

THE UNIVERSAL METHOD TO SUM SERIES FURTHER PROMOTED *

Leonhard Euler

§1 The universal Method to sum series, which I explained at the end of the last year, certainly extends very far, since from the given general term of the series alone it exhibits a formula equal to the sum of the series.; nevertheless it is difficult to accommodate it to series of such a kind, whose general terms cannot be expressed algebraically, but involve either exponential or even transcendental quantities. For, since having put the term, whose index is x , X the sum of the series from the first term until to X is equal to

$$\int Xdx + \frac{X}{1 \cdot 2} + \frac{dX}{1 \cdot 2 \cdot 3 \cdot 2dx} - \frac{d^3X}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6dx^3} + \text{etc.},$$

it is easily clear, if X involves at least quantities of this kind a^x , that so the expression $\int Xdx$ as the differentials of X are led to logarithms, whence a very large inconvenience arises in assigning the sum in question at least approximately.

§2 Furthermore, even if X was an algebraic quantity, nevertheless in many cases the differentials, which must be summed to obtain the sum, become so complicated that they can not only be exhibited in a difficult manner, but also yield a slowly converging series, as it happens in the series

$$\frac{1}{3} + \frac{1}{35} + \frac{1}{99} + \text{etc.}$$

*Original title: "Methodus universalis Series summandi ulterius promota", first published in „*Commentarii academiae scientiarum Petropolitanae* 8 (1736), 1741, p. 147-158", reprinted in in „*Opera Omnia: Series* 1, Volume 14, pp. 124 - 137 ", Eneström-Number E55, translated by: Alexander Aycock for „Euler-Kreis Mainz"

containing the quadrature of the circle. The main reason for this difficulty is that I assumed indices of the terms to grow by one unit; if I would have assumed them to grow by another number, a more tractable general term X might have been arisen. Finally, if the general term X cannot even be exhibited, as it happens in most series, then the given formula exhibiting the sum cannot even have any use.

§3 I have thought about this a long time how I could overcome these difficulties and observed that from the same principle, by means of which I had found that formula, also other formulas suitable to sum series can be found; having exhibited these for every series the formula is to be chosen, which is the most convenient. But from each class of formulas of this kind it seems to be convenient that two formulas are given, the one of which is apt to sum series from the first term up to a given term, of which kind the formula already mentioned was, but the other is apt to sum series from a given term to infinity. For, although this last summation follows from the first, it will nevertheless be useful to have given a peculiar formula for this case.

§4 Therefore, I will start from series, whose general term can be exhibited algebraically, for what also the method given in the preceding paper serves; but I will assume the indices to proceed in an arbitrary arithmetic progression, that the found formula extends further and will more often provide a convenient calculation. Therefore, let the series to be summed from the beginning to a given term with indices be the following

$$\begin{array}{ccccccc} a & a + b & a + 2b & & & & x \\ A & + B & + C & + \cdots & + X & = S, \end{array}$$

where the indices grow by the quantity b and the index of the first term A is a . Put the sum of this series $= S$; if in this expression for x one substitutes $x - b$, it is perspicuous that it will exhibit the same sum having taken away X or that it will be equal to $S - X$. But if in S instead of x one puts $x - b$, then it will arise

$$S - \frac{bdS}{1dx} + \frac{b^2ddS}{1 \cdot 2dx^2} - \frac{b^3d^3S}{1 \cdot 2 \cdot 3dx^3} + \text{etc.},$$

whence one will have the following equation

$$X = \frac{bdS}{1dx} - \frac{b^2ddS}{1 \cdot 2dx^2} + \frac{b^3d^3S}{1 \cdot 2 \cdot 3dx^3} - \frac{b^4d^4S}{1 \cdot 2 \cdot 3 \cdot 4dx^4} + \text{etc.}$$

But from this equation one finds this formula

$$S = \int \frac{Xdx}{b} + \frac{X}{1 \cdot 2} + \frac{bdX}{1 \cdot 2 \cdot 3 \cdot 2dx} - \frac{b^3d^3X}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6dx^3} + \frac{b^5d^5X}{1 \cdot 2 \cdot 3 \cdot \dots \cdot 7 \cdot 6dx^5}$$

$$- \frac{3b^7d^7X}{1 \cdot 2 \cdot 3 \cdot 9 \cdot 10dx^7} + \frac{5b^9d^9X}{1 \cdot 2 \cdot 3 \cdot \dots \cdot 11 \cdot 6dx^9} - \frac{691b^{11}d^{11}X}{1 \cdot 2 \cdot 3 \cdot \dots \cdot 12 \cdot 210dx^{11}}$$

$$+ \frac{35b^{13}d^{13}X}{1 \cdot 2 \cdot 3 \cdot \dots \cdot 15 \cdot 2dx^{13}} - \frac{3617b^{15}d^{15}X}{1 \cdot 2 \cdot 3 \cdot \dots \cdot 17 \cdot \dots \cdot 30dx^{15}} + \frac{43867b^{17}d^{17}X}{1 \cdot 2 \cdot 3 \cdot \dots \cdot 19 \cdot 42dx^{17}} - \text{etc.},$$

to which expression such a large constant is to be added that for $x = a$ it is $S = A$ or having put $x = a - b$ it is $S = 0$.

§5 If one puts $X = x^n$ or if the sum of this series is to be found

$$a^n + (a + b)^n + (a + 2b)^n + \dots + x^n,$$

it will be

$$\int Xdx = \frac{x^{n+1}}{n+1}, \quad \frac{dX}{dx} = nx^{n-1}, \quad \frac{d^3X}{dx^3} = n(n-1)(n-2)x^{n-3} \quad \text{etc.}$$

Hence the sum in question will be

$$S = \frac{x^{n+1}}{(n+1)b} + \frac{x^n}{1 \cdot 2} + \frac{nbx^{n-1}}{1 \cdot 2 \cdot 3 \cdot 2} - \frac{n(n-1)(n-2)b^3x^{n-3}}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6}$$

$$+ \frac{n(n-1)(n-2)(n-3)(n-4)b^5x^{n-5}}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 6} - \text{etc.} - \frac{a^{n+1}}{(n+1)b} + \frac{a^n}{1 \cdot 2} - \frac{nba^{n-1}}{1 \cdot 2 \cdot 3 \cdot 2}$$

$$+ \frac{n(n-1)(n-2)b^3a^{n-3}}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6} - \frac{n(n-1)(n-2)(n-3)(n-4)b^5a^{n-5}}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 6} + \text{etc.}$$

having added the corresponding constant. Therefore, the sum of the series

$$a + (a + b) + (a + 2b) + \dots + x$$

will be

$$= \frac{x^2}{2b} + \frac{x}{2} + \frac{b}{12} - \frac{a^2}{2b} + \frac{a}{2} - \frac{b}{12} = \frac{x^2 - a^2 + bx + ab}{2b}$$

and the sum of the series

$$\begin{aligned} a^2 + (a+b)^2 + (a+2b)^3 + \dots + x^2 &= \frac{x^3}{3b} + \frac{x^2}{2} + \frac{bx}{6} - \frac{a^3}{3b} + \frac{a^2}{2} - \frac{ab}{6} \\ &= \frac{2x^3 - 2a^3 + 3bx^2 + 3a^2b + b^2x - ab^2}{6b}. \end{aligned}$$

These expressions are similar to those, which I gave for the sum of powers of natural numbers in the superior dissertation, and from them they are also easily formed.

§6 Now to find the other formula of this class let a series to be summed from the first term X , whose index shall be x , to infinity, namely this one

$$\begin{array}{ccccccc} x & x+b & x+2b & & & & \\ X & + Y & + Z & + \text{etc. to infinity} & = & S. \end{array}$$

Therefore, if in the sum S for x one writes $x+b$, $S-X$ will arise; hence it will be

$$X = -\frac{bdS}{1dx} - \frac{b^2ddS}{1 \cdot 2dx^2} - \frac{b^3d^3S}{1 \cdot 2 \cdot 3dx^3} - \text{etc.},$$

whence as above it will be found

$$\begin{aligned} S = & - \int \frac{Xdx}{b} + \frac{X}{1 \cdot 2} + \frac{bdX}{1 \cdot 2 \cdot 3 \cdot 2dx} - \frac{b^3d^3X}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6dx^3} - \frac{b^5d^5X}{1 \cdot 2 \cdot 3 \cdot \dots \cdot 7 \cdot 6dx^5} \\ & + \frac{3b^7d^7X}{1 \cdot 2 \cdot 3 \cdot 9 \cdot 10dx^7} - \frac{5b^9d^9X}{1 \cdot 2 \cdot 3 \cdot \dots \cdot 11 \cdot 6dx^9} + \frac{691b^{11}d^{11}X}{1 \cdot 2 \cdot 3 \cdot \dots \cdot 12 \cdot 210dx^{11}} \\ & - \frac{35b^{13}d^{13}X}{1 \cdot 2 \cdot 3 \cdot \dots \cdot 15 \cdot 2dx^{13}} + \frac{3617b^{15}d^{15}X}{1 \cdot 2 \cdot 3 \cdot \dots \cdot 17 \cdot \dots \cdot 30dx^{15}} - \frac{43867b^{17}d^{17}X}{1 \cdot 2 \cdot 3 \cdot \dots \cdot 19 \cdot 42dx^{17}} + \text{etc.} \end{aligned}$$

To this formula such a large constant is to be added that it is $S=0$, if one puts $x=\infty$; for, if X already was the infinitesimal term or the last in the series, the sum must be vanishing, if the series has a finite sum, of course, to which case this formula is accommodated.

§7 That the use of this formula becomes clear, let $X = \frac{1}{x^2}$ or this series

$$\frac{1}{x^2} + \frac{1}{(x+b)^2} + \frac{1}{(x+2b)^2} + \text{etc.}$$

to be summed to infinity; therefore, because of

$$\int X dx = -\frac{1}{x}, \quad \frac{dX}{dx} = \frac{-2}{x^3}, \quad \frac{d^2X}{dx^2} = \frac{-2 \cdot 3 \cdot 4}{x^5} \quad \text{etc.}$$

it will be

$$S = \frac{1}{bx} + \frac{1}{2x^2} + \frac{b}{6x^3} - \frac{b^3}{3030x^5} + \frac{b^5}{42x^7} - \frac{b^7}{30x^9} + \frac{5b^9}{66x^{11}} - \frac{691b^{11}}{2730x^{13}} \\ + \frac{7b^{13}}{6x^{15}} - \frac{3617b^{15}}{510x^{17}} + \frac{43867b^{17}}{35910x^{19}} - \text{etc.},$$

which expression does not require a constant, since it vanishes per se having put $x = \infty$. But the series will converge the more the greater x was with respect to b . Hence, if several initial terms of the given series are actually added, the sum of the remaining ones found by this method added to that aggregate will give the sum of the propounded series continued to infinity.

§8 But having mentioned all these things, which only render the first rule more convenient, I proceed to the summation of series for which that formula does not suffice. For, let a series to be summed in which the signs of the terms alternate as

$$a \quad a+b \quad a+2b \quad a+3b \quad \dots \quad x \quad x+b \\ A \quad + B \quad + C \quad - D \quad + \dots + X \quad - Y = S;$$

since in this series Y is such a function of $x+b$ as X is one of x , it will be

$$Y = X + \frac{bdX}{1dx} + \frac{b^2ddX}{1 \cdot 2dx^2} + \frac{b^3d^3X}{1 \cdot 2 \cdot 3dx^3} + \text{etc.}$$

Further, if in S instead of x one puts $x-2b$, $S-X+Y$ will arise; therefore, it will be

$$-Y+X = \frac{2bdS}{1dx} - \frac{4b^2ddS}{1 \cdot 2dx^2} + \frac{8b^3d^3S}{1 \cdot 2 \cdot 3dx^3} - \frac{16b^4d^4S}{1 \cdot 2 \cdot 3 \cdot 4dx^4} + \frac{32b^5d^5S}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5dx^5} - \text{etc.}$$

$$= -\frac{bdX}{1dx} - \frac{b^2d^2X}{1 \cdot 2dx^2} - \frac{b^3d^3X}{1 \cdot 2 \cdot 3dx^3} - \frac{b^4d^4X}{1 \cdot 2 \cdot 3 \cdot 4dx^4} - \text{etc.}$$

Put

$$\frac{dS}{dx} = \frac{\alpha dX}{dx} + \frac{\beta d^2X}{dx^2} + \frac{\gamma d^3X}{dx^3} + \frac{\delta d^4X}{dx^4} + \text{etc.}$$

and by comparing the homologous terms it will arise

$$S = C - \frac{X}{2} - \frac{3bdX}{4dx} - \frac{b^2ddX}{2dx^2} - \frac{3b^3d^3X}{16dx^3} - \frac{b^4d^4X}{24dx^4} - \frac{9b^5d^5X}{1440dx^5} - \text{etc.}$$

and by introducing Y it will be

$$S = C - Y + \frac{X}{2} + \frac{bdX}{1 \cdot 2 \cdot 2dx} - \frac{b^3d^3X}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 2dx^3} + \frac{3b^5d^5X}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 2dx^5} - \text{etc.}$$

And hence it will be

$$\begin{aligned} & A - B + C - D + E - F + \dots + X \\ = & \text{Const.} + \frac{X}{2} + \frac{bdX}{1 \cdot 2 \cdot 2dx} - \frac{b^3d^3X}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 2dx^3} + \frac{3b^5d^5X}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 2dx^5} - \text{etc.}, \end{aligned}$$

where the constant as before must be of such a nature that this sum becomes = A having put $x = a$.

§9 But having continued the terms of this formula further and having put the sum of this series

$$\begin{array}{cccccccc} a & a+b & a+2b & a+3b & & & & x \\ A & -B & +C & -D & + & \dots & + & X = S; \end{array}$$

it will arise

$$\begin{aligned} S = & \text{Const.} + \frac{X}{1 \cdot 2} + \frac{bdX}{1 \cdot 2 \cdot 2dx} - \frac{b^3d^3X}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 2dx^3} + \frac{3b^5d^5X}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 2dx^5} \\ & - \frac{17b^7d^7X}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 8 \cdot 1dx^7} + \frac{155b^9d^9X}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 7 \cdot 8 \cdot 9 \cdot 10 \cdot 2dx^9} - \frac{2073b^{11}d^{11}X}{1 \cdot 2 \cdot 3 \dots 12 \cdot 2dx^{11}} \end{aligned}$$

$$+ \frac{38227b^{13}d^{13}X}{1 \cdot 2 \cdot 3 \dots 14 \cdot 2dx^{13}} - \text{etc.}$$

Therefore, if the value of this progression is to be found

$$a^2 - (a + b)^2 + (a + 2b)^2 - (a + 3b)^2 + \dots + x^2,$$

which shall be S , because of

$$\frac{dX}{dx} = 2x$$

it will be

$$S = \text{Const.} + \frac{x^2}{2} + \frac{bx}{2}.$$

But the constant C on the other will be found by putting $x = a$ and it will be

$$S = \frac{a^2 - ab + x^2 + bx}{2}$$

For the sake of an example, it will be

$$1 - 4 + 9 - 16 + 25 - \dots + 121 = 66.$$

§10 Now let us consider infinite series of this kind, namely

$$S = X - Y + Z - \text{etc. to infinity}$$

Therefore, it will be

$$S + \frac{2bdS}{1dx} + \frac{4b^2ddS}{1 \cdot 2dx^2} + \frac{8b^3d^3S}{1 \cdot 2 \cdot 3dx^3} + \text{etc.} = S - X + Y$$

or

$$X - Y = -\frac{2bdS}{1dx} - \frac{4b^2ddS}{1 \cdot 2dx^2} - \text{etc.}$$

whence, because it is

$$Y = X + \frac{bdX}{1dx} + \frac{b^2ddX}{1 \cdot 2dx^2} + \frac{b^3d^3X}{1 \cdot 2 \cdot 3dx^3} + \text{etc.}$$

it will be found

$$S = \frac{X}{1 \cdot 2} - \frac{bdX}{1 \cdot 2 \cdot 2dx} + \frac{b^3 d^3 X}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 2dx^3} - \frac{3b^5 d^5 X}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5 \cdot 6 \cdot 2dx^5} + \frac{17b^7 d^7 X}{1 \cdot 2 \cdot 3 \cdots 8 \cdot 2dx^7} \\ - \frac{155b^9 d^9 X}{1 \cdot 2 \cdot 3 \cdots 10 \cdot 2dx^9} + \frac{2073b^{11} d^{11} X}{1 \cdot 2 \cdot 3 \cdots 12 \cdot 2dx^{11}} - \frac{38227b^{13} d^{13} X}{1 \cdot 2 \cdot 3 \cdots 14 \cdot 2dx^{13}} + \text{etc.} + \text{Const.}$$

This constant must be of such a nature that it is $S = 0$ for $x = \infty$. Therefore, by means of this formula many other slowly converging series, in which the signs of the terms alternate, can be summed approximately.

§11 Let $X = \frac{1}{x}$ such that the sum of this series must be found

$$\frac{1}{x} - \frac{1}{x+b} + \frac{1}{x+2b} - \frac{1}{x+3b} + \text{etc. to infinity}$$

Therefore, because it is $X = \frac{1}{x}$, it will be

$$\frac{dX}{dx} = \frac{-1}{x^2}, \quad \frac{d^3 X}{dx^3} = \frac{-2 \cdot 3}{x^4}, \quad \frac{d^5 X}{dx^5} = \frac{-2 \cdot 3 \cdot 4 \cdot 5}{x^6} \quad \text{etc.}$$

and hence it will be

$$S = \frac{1}{2x} + \frac{b}{2 \cdot 2x^2} - \frac{b^3}{4 \cdot 2x^4} + \frac{3b^5}{6 \cdot 2x^6} - \frac{17b^7}{8 \cdot 2x^8} + \frac{155b^9}{10 \cdot 2x^{10}} - \frac{203b^{11}}{12 \cdot 2x^{12}} + \frac{38227b^{13}}{14 \cdot 2x^{14}} - \text{etc.},$$

where the addition of the constant is not necessary. For the sake of an example let $b = 2$ and $x = 25$; the sum of the series

$$\frac{1}{25} - \frac{1}{27} + \frac{1}{29} - \frac{1}{31} + \frac{1}{33} - \text{etc.}$$

will be

$$= \frac{2}{100} + \frac{8}{100^2} - \frac{256}{100^4} + \frac{8 \cdot 4^6}{100^6} - \frac{17 \cdot 8 \cdot 4^8}{100^8} + \frac{31 \cdot 128 \cdot 4^{10}}{100^{10}} - \frac{691 \cdot 256 \cdot 4^{12}}{100^{12}} \\ + \frac{5461 \cdot 2048 \cdot 4^{14}}{100^{14}} - \text{etc.} = 0.020797471915 \quad \text{approximately;}$$

but if to the sum of the preceding terms $1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \cdots - \frac{1}{23}$ the found sum is actually added, the fourth part of the circumference of this circle will arise, while the diameter of that circle is = 1.

§12 Now, the formulas given until now require, that the calculation can be performed easily, that X is an algebraic function of x , whose differentials of each degree can be exhibited in a convenient matter. For, these formulas could hardly or not even hardly be used, if an exponential quantity of this kind n^x would be in the general term of the progression. Therefore, it will be convenient to find peculiar summation formulas for progressions contained in this general term Xn^x , where X as before denotes an algebraic function of x . Therefore, let this series

$$An^a + Bn^{a+b} + Cn^{a+2b} + \dots + Xn^x = S;$$

be propounded to be summed and the sum be put $= Sn^x$. But this formula having put $x - b$ instead of x will go over into this one

$$n^{x-b} \left(S - \frac{bdS}{1dx} + \frac{b^2d^2S}{1 \cdot 2dx^2} - \frac{b^3d^3S}{1 \cdot 2 \cdot 3dx^3} + \text{etc.} \right),$$

which must be equal to the first sum Sn^x having taken away the last term Xn^x . Therefore, one will have this equation

$$Sn^b - Xn^b = S - \frac{bdS}{1dx} + \frac{b^2d^2S}{1 \cdot 2dx^2} - \frac{b^3d^3S}{1 \cdot 2 \cdot 3dx^3} + \frac{b^4d^4S}{1 \cdot 2 \cdot 3 \cdot 4dx^4} - \text{etc.}$$

From this equation the value of S must be found.

§13 Therefore, put $n^b = m$ and it will be

$$S = \frac{mX}{m-1} - \frac{\alpha bdX}{1(m-1)^2dx} + \frac{\beta b^2 ddX}{1 \cdot 2(m-1)^3dx^3} - \frac{\gamma b^3 d^3 X}{1 \cdot 2 \cdot 3(m-1)^4dx^3} \\ + \frac{\delta b^4 d^4 X}{1 \cdot 2 \cdot 3 \cdot 4(m-1)^5dx^4} - \frac{\epsilon b^5 d^5 X}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5(m-1)^6dx^5} + \text{etc.}$$

Hence by comparing the homologous terms having for the sake of brevity put $m - 1 = p$ it will be as follows

$$\begin{aligned}
\alpha &= m, \\
\beta &= 2\alpha + mp, \\
\gamma &= 3\beta + 3\alpha p + mp^2, \\
\delta &= 4\gamma + 6\beta p + 4\alpha p^2 + mp^3, \\
\varepsilon &= 5\delta + 10\gamma p + 10\beta p^2 + 5\alpha p^3 + mp^4
\end{aligned}$$

whence for the letters $\alpha, \beta, \gamma, \delta$ etc. the following values are obtained

$$\begin{aligned}
\alpha &= m, \\
\beta &= 2m + mp, \\
\gamma &= 6m + 6mp + mp^2, \\
\delta &= 24m + 36mp + 14mp^2 + mp^3, \\
\varepsilon &= 120m + 240mp + 150mp^2 + 30mp^3 + mp^4
\end{aligned}$$

etc.

Here, each number is a multiple of the one written above combined with the preceding one, as

$$\begin{aligned}
30 &= 2(1 + 11), \\
150 &= 3(14 + 36), \\
240 &= 4(36 + 24), \\
120 &= 5(24 + 0),
\end{aligned}$$

or having resubstituted $m - 1$ instead of p it is as follows

1. $\alpha = m,$
2. $\beta = m + m^2,$
3. $\gamma = m + 4m^2 + m^3,$
4. $\delta = m + 11m^2 + 11m^3 + m^4,$
5. $\varepsilon = m + 26m^2 + 66m^3 + 26m^4 + m^5,$
6. $\zeta = m + 57m^2 + 302m^3 + 302m^4 + 57m^5 + m^6,$

etc.,

which values proceed in such a way that the coefficient ψ , whose index is k , is

$$\begin{aligned}
&= m + \left(2^k - \frac{k+1}{1}\right) m^2 + \left(3^k - \frac{k+1}{1}2^k + \frac{(k+1)k}{1 \cdot 2}\right) m^3 \\
&+ \left(4^k - \frac{k+1}{1}3^k + \frac{(k+1)k}{1 \cdot 2}2^k - \frac{(k+1)k(k-1)}{1 \cdot 2 \cdot 3}\right) m^4 + \dots + m^k.
\end{aligned}$$

§14 Therefore, from these the sum of the propounded series

$$\begin{array}{cccc}
a & a+b & a+2b & x \\
An^a & + Bn^{a+b} & + Cn^{a+2b} & + \dots + Xn^x
\end{array}$$

is calculated

$$= n^x \left\{ \begin{array}{l} \frac{n^b X}{n^b - 1} - \frac{n^b b dX}{1(n^b - 1)dx} + \frac{(n^{2b} + n^b)^2 b^2 d dX}{1 \cdot 2(n^b - 1)^3 dx^2} - \frac{(n^{3b} + 4n^{2b} + n^b) b^3 d^3 X}{1 \cdot 2 \cdot 3(n^b - 1)^4 dx^3} \\ + \frac{(n^{4b} + 11n^{3b} + 11n^{2b} + n^b) b^4 d^4 X}{1 \cdot 2 \cdot 3 \cdot 4(n^b - 1)^5 dx^4} - \text{etc.} + \text{Const.}, \end{array} \right\} + \text{Const.},$$

which constant must be of such a nature that having put $x = a$ the sum is $= An^a$. If one puts $n^b = -1$, the series will go over into a pure algebraic one, in which the signs alternate, and hence by putting -1 instead of n^b the same formula we found for the same case already above in § 9 results. But the sum of the series running to infinity

$$\begin{array}{cccc}
x & x+b & x+2b & \\
An^x & + Yn^{x+b} & + Zn^{x+2b} & + \text{etc. to infinity}
\end{array}$$

is

$$= n^x \left\{ \begin{array}{l} \frac{-X}{n^b - 1} + \frac{n^b b dX}{1(n^b - 1)^2 dx} - \frac{(n^{2b} + n^b) b^2 d dX}{1 \cdot 2(n^b - 1)^3 dx^2} + \frac{(n^{3b} + 4n^{2b} + n^b) b^3 d^3 X}{1 \cdot 2 \cdot 3(n^b - 1)^4 dx^3} \\ - \frac{(n^{4b} + 11n^{3b} + 11n^{2b} + n^b) b^4 d^4 X}{1 \cdot 2 \cdot 3 \cdot 4(n^b - 1)^5 dx^4} + \text{etc.} + \text{Const.}, \end{array} \right\} + \text{Const.},$$

which constant must be of such a nature that having put $x = \infty$ the sum becomes $= 0$; this always happens per se such that a constant is not necessary, if the series converges and has a finite sum, of course.

§15 From the formula for summing series of this kind up to a given term it is understood, if X was such an algebraic function of x , that finally its higher differentials vanish, then the summatory term can indeed be exhibited. Therefore, if the series, whose general term is X , was summable, then also the series, whose general term is Xn^x , will be summable. So having propounded the series

$$a^2n^a + (a+b)n^{a+b} + (a+2b)n^{a+2b} + \dots + x^2n^x$$

its sum will be

$$= n^x \left(\frac{n^b x^2}{n^b - 1} - \frac{2n^b x}{(n^b - 1)^2} + \frac{(n^{2b} + n^b)b^2}{(n^b - 1)^3} \right) + \text{Const.},$$

which constant having put $x = a$ and the sum $= a^2n^a$ it will arise as

$$n^a \left(a^2 - \frac{n^b a^2}{n^b - 1} + \frac{2n^b ab}{(n^b - 1)^2} - \frac{(n^{2b} + n^b)b^2}{(n^b - 1)^3} \right).$$

§16 Furthermore, the other formula has an immense use in the summation of infinite series; that this becomes more clear, let this series be propounded

$$\frac{n^x}{x} + \frac{n^{x+2}}{x+2} + \frac{n^{x+4}}{x+4} + \frac{n^{x+6}}{x+6} + \text{etc.},$$

whose sum shall be $= S$. Therefore, it will be $b = 2$ and $X = \frac{1}{x}$, whence the sum will become

$$S = n^x \left\{ \begin{array}{l} \frac{-1}{(n^2 - 1)x} - \frac{2n^2}{(n^2 - 1)^2 x^2} - \frac{4(n^4 + n^2)}{(n^2 - 1)^3 x^3} - \frac{8(n^6 + 4n^4 + n^2)}{(n^2 - 1)^4 x^4} \\ - \frac{16(n^8 + 11n^6 + 11n^4 + n^2)}{(n^2 - 1)^5 x^5} - \text{etc.} \end{array} \right\}.$$

Now let $x = 25$ and $n^2 = -\frac{1}{3}$ or $n = \frac{1}{\sqrt{-3}}$; it will be

$$\begin{aligned} & \frac{1}{\sqrt{-3}} \left(\frac{1}{25 \cdot 3^{12}} - \frac{1}{27 \cdot 3^{15}} + \frac{1}{29 \cdot 3^{14}} - \frac{1}{31 \cdot 3^{15}} + \text{etc.} \right) \\ &= \frac{1}{3^{12} \sqrt{-3}} \left(\frac{3}{4 \cdot 25} + \frac{3}{8 \cdot 25^2} - \frac{3}{8 \cdot 25^3} - \frac{3}{16 \cdot 25^4} + \frac{15}{8 \cdot 25^5} + \text{etc.} \right). \end{aligned}$$

Therefore, because in the circle with radius = 1 the arc of thirty degree is

$$= \frac{1}{\sqrt{3}} \left(1 - \frac{1}{3 \cdot 3} + \frac{1}{5 \cdot 3^2} - \frac{1}{7 \cdot 3^3} + \frac{1}{9 \cdot 3^4} - \text{etc.} \right)$$

if 12 of these terms are actually added, the following remaining ones will be

$$\begin{aligned} & \frac{1}{\sqrt{3}} \left(\frac{1}{25 \cdot 3^{12}} - \frac{1}{27 \cdot 3^{15}} + \frac{1}{29 \cdot 3^{14}} - \text{etc.} \right) \\ &= \frac{\sqrt{3}}{3^{12}} \left(\frac{1}{4 \cdot 25} + \frac{1}{8 \cdot 25^2} - \frac{1}{8 \cdot 25^3} - \frac{1}{16 \cdot 25^4} + \frac{55}{8 \cdot 25^5} + \text{etc.} \right). \end{aligned}$$

§17 If the terms of the series to be summed were composed of factors such that the series has a form of this kind

$$a \quad a + b \quad a + 2b \quad \dots \quad x \\ A + AB + ABC + \dots + ABC \dots VX,$$

put the sum

$$= S \cdot ABC \dots VX.$$

Therefore, this sum having put $x - b$ instead of x will go over into this one

$$ABC \dots V \left(S - \frac{bdS}{1dx} + \frac{b^2ddS}{1 \cdot 2dx^2} - \frac{b^3d^3S}{1 \cdot 2 \cdot 3dx^3} + \text{etc.} \right),$$

which must be equal to the first term having taken away the last term, this means the quantity

$$AB \cdot V(SX - X).$$

Therefore, one will have this equation

$$SX - X = S - \frac{bdS}{1dx} + \frac{b^2ddS}{1 \cdot 2dx^2} - \frac{b^3d^3S}{1 \cdot 2 \cdot 3dx^3} + \frac{b^4d^4S}{1 \cdot 2 \cdot 3 \cdot 4dx^4} - \text{etc.}$$

From this equation the found value of S will be this

$$S = \frac{X}{X - 1} + E + F + G + \text{etc.},$$

which terms proceed in such a way that having put

$$\frac{x}{X-1} = D \quad (1)$$

it is

$$\begin{aligned} E &= -\frac{bdD}{(X-1)dx'} \\ F &= -\frac{bdE}{(X-1)dx} + \frac{b^2ddD}{1 \cdot 2(X-1)dx^2'} \\ G &= -\frac{bdF}{(X-1)dx} + \frac{b^2ddE}{1 \cdot 2(X-1)dx^2} - \frac{b^3d^3D}{1 \cdot 2 \cdot 3(X-1)dx^3} \end{aligned}$$

and so forth, that hence the sum of the propounded series is

$$= ABC \cdots VX(D + E + F + G + \text{etc.}).$$

But the sum of the series continued to infinity

$$ABC \cdots VX + ABCD \cdots VXY + \text{etc.}$$

will be

$$ABC \cdots VX(1 - D - E - F - G - \text{etc.}) + \text{Const.}$$

As if the sum of this series is in question

$$\frac{1}{1 \cdot 2 \cdot 3 \cdots x} + \frac{1}{1 \cdot 2 \cdot 3 \cdots x(x+1)} + \text{etc. to infinity,}$$

it will be

$$b = 1, \quad X = \frac{1}{x} \quad \text{and} \quad D = \frac{1}{1-x},$$

but then

$$E = \frac{-x}{(1-x)^3}, \quad F = \frac{x(2+x)}{(1-x)^5} \quad \text{etc.};$$

therefore, the sum of the propounded series because of the vanishing constant will be

$$= \frac{1}{1 \cdot 2 \cdot 3 \cdots x} \left(1 + \frac{1}{x-1} - \frac{x}{(x-1)^3} + \frac{x(x+2)}{(x-1)^5} - \text{etc.} \right).$$

From these things given it is indeed easily understood, a form of what kind of the sum must be assumed in each case that the sum is found by minimal work.