

# ON THE INTEGRATION OF DIFFERENTIAL EQUATIONS \*

Leonhard Euler

§1 Here, I consider differential equations of first degree, which involve only two variables, which therefore can be represented in this general form

$$Mdx + Ndy = 0,$$

if  $M$  and  $N$  denote any arbitrary functions of the two variables  $x$  and  $y$ . But it was proved that an equation of this kind always expresses a certain relation between the variables  $x$  and  $y$ , by which for each value of the one the values for the other are defined. But because this finite relation between the variables must be found by means of integration, the integral equation, if it is extended to its complete generality, will receive a constant quantity, which, while it is completely arbitrary, contains infinitely many integral equations, which all satisfy the differential equation equally.

§2 Therefore, having propounded any differential equation of this kind

$$Mdx + Ndy = 0$$

the whole task of the Analysis consists in this that a finite equation between the same variables  $x$  and  $y$  is found, which expresses the same among those relations as the differential equation, and this in the broadest sense, of course

---

\*Original title: " De integratione aequationum differentialium ", first published in „*Novi Commentarii academiae scientiarum Petropolitanae* 8, (1760/1), 1763, p.3-63", reprinted in in „*Opera Omnia*: Series 1, Volume 22, pp. 334 - 394 ", Eneström-Number E269, translated by: Alexander Aycock for „Euler-Kreis Mainz"

such that it contains a certain arbitrary constant, which is not in the differential equation. But if this question is propounded in this most general form, until now no way was found to get to its solution; and all cases, which could be solved until now, can be reduced to a tiny number, such that in this part of Analysis, as in the remaining ones, still very large increments are desired; and therefore, the complete cognition of all secrets of this science can never be expected.

§3 Almost everything achieved in this task until now can be referred to these cases, in which the differential equation

$$Mdx + Ndy = 0$$

either immediately admits a separation of variables, or by means of suitable substitutions can be reduced to such a form. For, if by introducing two new variables  $v$  and  $z$  instead of  $x$  and  $y$ , the propounded differential equation can be transformed in a form of this kind

$$Vdv + Zdz = 0,$$

in which  $V$  is a function only of  $v$  and  $Z$  only of  $z$ , the whole task will be completed, while the complete integral equation will be:

$$\int Vdv + \int Zdz = \text{Const.},$$

which manifestly contains that arbitrary constant introduced by the general integration. And to this almost all artifices reduce, which the Analysts have used in the resolution of equations of this kind.

§4 Therefore, if the propounded differential equation does not immediately admit the separation of variables, the whole task used to be that appropriate substitutions, which open the way to the separation, are investigated, where often the highest ingeniousness, which the Geometers used to reach his goal, must be admired. Since there is nevertheless no certain way known to investigate substitutions of this kind, this method seems rather unnatural, whence I decided, to consider a maybe not new, but nevertheless still not sufficiently developed method in more detail, since which does not require substitutions seems to be more natural being found on the nature of differentials and even contains the first method, at least partially, in it as a special case.

§5 Having reduced the differential equation to this form

$$Mdx + Ndy = 0$$

consider the formula  $Mdx + Ndy$  without taking into account that it has to vanish, and examine, whether it is a differential of a certain function of  $x$  and  $y$  or not. This examination is to be done in such a way, as it was explained already abundantly in different sources; of course, both functions  $M$  and  $N$  must be differentiated, and since their differentials must have a form of this kind

$$dM = pdx + qdy \quad \text{and} \quad dN = rdx + sdy,$$

see, whether it is  $q = r$  or not. For, if it was  $q = r$ , this is an infallible criterion that the formula  $Mdx + Ndy$  is integrable: But if it was not  $q = r$ , it is equally certain, that this formula did not arise from a differentiation of a certain finite function of  $x$  and  $y$ . From this the whole question is reduced to two cases, of which the one holds, if it was  $q = r$ , the other, if these two quantities  $q$  and  $r$  were not equal to each other.

§6 Therefore, to see the equality, or inequality, of the quantities  $q$  and  $r$ , it is not even necessary that the functions  $M$  and  $N$  are expanded completely by differentiation, but it suffices in the function  $M$ , which is connected to  $dx$ , to consider the quantity  $x$  as constant, and only ask for its partial differential, which is obtained from the variability of  $y$  only, since this way the term  $qdy$  is obtained, but I usually denote the value of  $q$  found this way by the sign  $\left(\frac{dM}{dy}\right)$ . In similar manner, differentiate the other function  $N$ , which is connected to  $dy$ , in such a way, that  $y$  is treated as a constant, and from the variability of  $x$  only the part  $rdx$  is obtained, where I equally express the value of  $r$  by  $\left(\frac{dN}{dx}\right)$ . Therefore, if the formula  $Mdx + Ndy$  was of such a nature that it is

$$\left(\frac{dM}{dy}\right) = \left(\frac{dN}{dx}\right),$$

it is integrable, and its integral can be found the following way. Having done this, if the condition of criterion is not satisfied, let us see how it is to be proceeded.

## PROBLEM 1

§7 If the differential equation

$$Mdx + Ndy = 0$$

was of such a nature that it is

$$\left(\frac{dM}{dy}\right) = \left(\frac{dN}{dx}\right),$$

to find its integral equation.

## SOLUTION

If it was

$$\left(\frac{dM}{dy}\right) = \left(\frac{dN}{dx}\right),$$

then a finite function of the two variables  $x$  and  $y$  is given, which differentiated yields  $Mdx + Ndy$ . Let  $V$  be this function, and because it is

$$dV = Mdx + Ndy,$$

$Mdx$  will be the differential of  $V$ , if only  $x$  is taken as a variable, and  $Ndy$  its differential, if only  $y$  is taken as a variable. Therefore, hence vice versa  $V$  will be found, if either  $Mdx$  is integrated, having considered  $y$  as constant, or  $Ndy$  is integrated, having considered  $x$  as constant: And so this operation is reduced to the integration of a differential formula involving only one variable which is postulated to be possible in this case, might it succeed algebraically or require the quadrature of curves. But because reasoning as this the quantity  $V$  is found in two ways, and the one integration instead of the constant leads to an arbitrary function of  $y$ , the other on the other hand assumes a function of  $x$ , such that it is

$$\text{both } V = \int Mdx + Y \quad \text{and} \quad V = \int Ndy + X,$$

it is always possible to define these functions  $Y$  of  $y$  and  $X$  of  $x$  in such a way that it is  $\int Mdx + Y = \int Ndy + X$ , which is easily achieved in each case. Since having done this the quantity  $V$  is the integral of the formula  $Mdx + Ndy$ , it is evident that the integral equation of the propounded equation  $Mdx + Ndy = 0$

will be  $V = \text{Const.}$ , and it will be the complete integral equation, since it involves an arbitrary constant.

#### COROLLARY 1

§8 In this case the case of separated equations is immediately contained. For, if  $M$  was a function of  $x$  only, and  $N$  a function of  $y$  only, it will certainly be

$$\left(\frac{dM}{dy}\right) = 0 \quad \text{and} \quad \left(\frac{dN}{dx}\right) = 0 \quad \text{and hence} \quad \left(\frac{dM}{dy}\right) = \left(\frac{dN}{dx}\right);$$

therefore, this is the simplest case, which the problem contains in it.

#### COROLLARY 2

§9 But if in the differential equation

$$Mdx + Ndy = 0$$

$M$  was a function of  $x$  only, and  $N$  of of  $y$  only, each of both parts is integrable separately, and the integral equation will be:

$$\int Mdx + \int Ndy = \text{Const.}$$

#### COROLLARY 3

§10 Furthermore, our problem provides us with the solution of infinitely many other differential equations, the common character of all of which consists in this that it is

$$\left(\frac{dM}{dy}\right) = \left(\frac{dN}{dx}\right),$$

and their resolution can be done by means of integration of formulas containing one single variable.

## SCHOLIUM 1

§11 Therefore, as often as in the differential equation  $Mdx + Ndy = 0$  it was  $\left(\frac{dM}{dy}\right) = \left(\frac{dN}{dx}\right)$ , its resolution has no difficulty, as long as the integration of formulas involving one variable is conceded; this can certainly justly be postulated. Nevertheless, the determination of those functions  $X$  and  $Y$  which have to be introduced instead of the constants, could seem to create certain inconveniences; but it will soon be found to vanish in the single cases. But to contract this operation even further, the second integration is not even necessary. For, after the one part  $Mdx$ , having considered  $y$  as constant, was integrated, which integral shall be  $= Q$ , set

$$V = Q + Y,$$

having meanwhile put  $Y$  for a indefinite function of  $y$ , which the other variable  $x$  does not go into at all. Then differentiate this quantity  $Q + Y$  again by treating  $x$  as a constant, and since the differential has to arise as  $= Ndy$ , from this condition the function  $Y$  will be determined most easily, since from the nature of the procedure itself hence the quantity  $x$  will be eliminated immediately. But having found this function  $Y$ , the integral equation will be  $Q + Y = \text{Const.}$ , which operation will be conveniently illustrated in the following examples.

## EXAMPLE 1

§12 To integrate this differential equation:

$$2axydx + axxdy - y^3dx - 3xyydy = 0.$$

Having compared this equation to the form  $Mdx + Ndy = 0$ , it will be:

$$M = 2axy - y^3 \quad \text{and} \quad N = axx - 3xyy.$$

Therefore, first it is to be checked, whether this case is contained in the problem, for which aim we want to find the values:

$$\left(\frac{dM}{dy}\right) = 2ax - 3yy \quad \text{and} \quad \left(\frac{dN}{dx}\right) = 2ax - 3yy,$$

because which are equal, the prescribed operation will necessarily succeed. But, haven taken  $y$  to be constant, one will find:

$$\int Mdx = axxy - y^3x + Y;$$

if the differential of this form is taken, having put  $x$  to be constant, it will arise:

$$axxdy - 3yyxdy + dY = Ndy,$$

and having resubstituted its value  $axx - 3xyy$  for for  $N$ , it will be  $dY = 0$ , whence  $Y = 0$  arises, or  $Y = \text{Const.}$ . Hence one will have the integral equation is question:

$$axxy - y^3x = \text{Const.}$$

### EXAMPLE 2

**§13** To integrate this differential equation:

$$\frac{ydy + xdx - 2ydx}{(y-x)^2} = 0.$$

Having compared this equation to the form  $Mdx + Ndy = 0$ , it will be:

$$M = \frac{x-2y}{(y-x)^2} \quad \text{and} \quad N = \frac{y}{(y-x)^2}.$$

Now, that it becomes plain, whether this equation is contained in the case of the problem, find the differential values:

$$\left(\frac{dM}{dy}\right) = \frac{2y}{(y-x)^2} \quad \text{and} \quad \left(\frac{dN}{dx}\right) = \frac{2y}{(y-x)^2},$$

since which are equal, the task will be successful. Hence according to the rule one concludes, having taken  $y$  to be constant, the integral:

$$\int Mdx = \int \frac{xdx - 2ydx}{(y-x)^2} = - \int \frac{dx}{y-x} - \int \frac{ydx}{(y-x)^2}$$

and one will find:

$$\int Mdx = \log(y-x) - \frac{y}{y-x} + Y,$$

whose differential, having taken  $x$  to be constant, must produce the other part  $Ndy$  of the propounded equation; hence one will have:

$$Ndy = \frac{dy}{y-x} + \frac{xdy}{(y-x)^2} + dY = \frac{ydy}{(y-x)^2} + dY.$$

Therefore, because it is

$$Ndy = \frac{ydy}{(y-x)^2}, \quad \text{it will } dY = 0 \quad \text{and } Y = 0,$$

for, the constant  $Y$  can be neglected, since it is introduced already in the integral equation, which will be:

$$\log(y-x) - \frac{y}{y-x} = \text{Const.}$$

### EXAMPLE 3

§14 To integrate this differential equation:

$$\frac{dx}{x} + \frac{yydx}{x^3} - \frac{ydy}{xx} + \frac{(ydx - xdy)\sqrt{xx+yy}}{x^3} = 0.$$

Having compared this equation to the from  $Mdx + Ndy = 0$ , we will have:

$$M = \frac{xx + yy + y\sqrt{xx+yy}}{x^3} \quad \text{and} \quad N = \frac{-y - \sqrt{xx+yy}}{xx},$$

whence for the exploration of the criterion find:

$$\left(\frac{dM}{dy}\right) = \frac{2y}{x^3} + \frac{\sqrt{xx+yy}}{x^3} + \frac{yy}{x^3\sqrt{xx+yy}}$$

and

$$\left(\frac{dN}{dx}\right) = \frac{2y}{x^3} + \frac{2\sqrt{xx+yy}}{x^3} - \frac{x}{xx\sqrt{xx+yy}},$$

since which values reduced become equal to each other, of course

$$\left(\frac{dM}{dy}\right) = \left(\frac{dN}{dx}\right) = \frac{2y}{x^3} + \frac{xx + 2yy}{x^3\sqrt{xx+yy}},$$

the resolution will be possible. Therefore, investigate, having taken  $y$  to be constant:



$$\int Mdx = \log x - \frac{yy}{2xx} + y \int \frac{dx}{x^3} \sqrt{xx + yy}.$$

But by means of the rules to integrate formulas involving one variable, since here  $y$  is considered to be constant, one finds:

$$\int \frac{ydx}{x^3} \sqrt{xx + yy} = \frac{-y\sqrt{xx + yy}}{2xx} + \frac{1}{2} \log \frac{\sqrt{xx + yy} - y}{y},$$

such that it is:

$$\int Mdx = \log x - \frac{yy}{2xx} - \frac{y\sqrt{xx + yy}}{2xx} + \frac{1}{2} \log \frac{\sqrt{xx + yy} - y}{y} + Y.$$

But since the differential of this quantity, having assumed  $x$  to be constant, has to yield

$$Ndy = -\frac{-ydy - dy\sqrt{xx + yy}}{xx},$$

we will obtain:

$$Ndy = \frac{-ydy}{xx} - \frac{dy\sqrt{xx + yy}}{2xx} - \frac{yydy}{2xx\sqrt{xx + yy}} - \frac{dy}{2y} - \frac{dy}{2\sqrt{xx + yy}} + dY,$$

having compared this form to that one it will be:

$$dY = -\frac{dy\sqrt{xx + yy}}{2xx} + \frac{yydy}{2xx\sqrt{xx + yy}} + \frac{dy}{2y} + \frac{dy}{2\sqrt{xx + yy}},$$

where the terms, which still contain  $x$ , cancel each other, such that it is

$$dY = \frac{dy}{2y} \quad \text{and} \quad Y = \frac{1}{2} \log y.$$

Having found this value for  $Y$ , one will obtain the integral equation in question:

$$\log x - \frac{yy}{2xx} - \frac{y\sqrt{xx + yy}}{2xx} + \frac{1}{2} \log(\sqrt{xx + yy} - y) = \text{Const.}$$

## SCHOLIUM 2

§15 From these examples it is sufficiently understood, how the prescribed operation is to be done, such that hence no difficulty causes any further inconveniences, except those always remaining from the integration of formulas involving one variable, when the integration cannot be done algebraically and does not allow to be reduced to the quadrature of the circle or hyperbola. But then the superior quadratures have to be treated in the same manner, and if these difficulties remain, they are not to be ascribed to this method. Therefore, it is possible to assume here, as often as the differential equation

$$Mdx + Ndy = 0$$

was of such a nature that in it it is

$$\left(\frac{dM}{dy}\right) = \left(\frac{dN}{dx}\right),$$

that so often the integration is possible for us; hence, I proceed to equations, in which this criterion is not satisfied.

### THEOREM

§16 If in the differential equation

$$Mdx + Ndy = 0$$

it was not

$$\left(\frac{dM}{dx}\right) = \left(\frac{dN}{dy}\right),$$

always a multiplier is given, multiplied by which the formula  $Mdx + Ndy$  becomes integrable.

### PROOF

Since it is not

$$\left(\frac{dM}{dy}\right) = \left(\frac{dN}{dx}\right),$$

the formula  $Mdx + Ndy$  will also not be integrable, or no function of  $x$  and  $y$  exists, whose differential is  $Mdx + Ndy$ . But here not the integral so of the formula  $Mdx + Ndy$  as of the equation  $Mdx + Ndy = 0$  is in question; and because the same equation holds, if it is multiplied by any function  $L$  of  $x$  and  $y$ , such that it is

$$LMdx + LNdy = 0,$$

it is to be demonstrated that always a function  $L$  of such a kind is given that the formula

$$LMdx + LNdy$$

becomes integrable. For, that this happens, it is necessary that it is:

$$\left(\frac{d.LM}{dy}\right) = \left(\frac{d.LN}{dx}\right),$$

or if one puts

$$dL = Pdx + Qdy,$$

because it is

$$\left(\frac{dL}{dy}\right) = Q \quad \text{and} \quad \left(\frac{dL}{dx}\right) = P,$$

the function  $L$  has to be of such a nature that it is:

$$L\left(\frac{dM}{dy}\right) + MQ = L\left(\frac{dN}{dx}\right) + NP.$$

But it is evident that this condition suffices to define the function  $L$ , if the formula  $Mdx + Ndy$  is multiplied by it, becomes integrable.

#### COROLLARY 1

§17 Therefore, having found the multiplier  $L$ , which renders the formula

$$Mdx + Ndy$$

integrable, the equation  $Mdx + Ndy = 0$  brought into the form

$$LMdx + LNdy = 0,$$

can be integrated by the method explained in the preceding problem.

### COROLLARY 2

§18 Having considered  $y$  as constant let the integral  $\int LMdx$  be in question, to which add such a function  $Y$  of  $y$  that, if the aggregate

$$\int LMdx + Y$$

is differentiated again, having considered  $x$  as constant now,  $LNdy$  arises. Having done this the integral equation will be

$$\int LMdx + Y = \text{Const.}$$

### COROLLARY 3

§19 Therefore, the multiplier must be of such a nature that having put

$$dL = Pdx + Qdy,$$

this equation is satisfied:

$$L \left( \frac{dM}{dy} \right) + MQ = L \left( \frac{dN}{dx} \right) + NP$$

or this one:

$$\frac{NP - MQ}{L} = \left( \frac{dM}{dy} \right) - \left( \frac{dN}{dx} \right),$$

whence it is manifest, if it would be

$$\left( \frac{dM}{dy} \right) = \left( \frac{dN}{dx} \right),$$

that for  $L$  one can take the unity, or any constant quantity, while it is  $P = 0$  and  $Q = 0$ .

## SCHOLIUM

§20 Therefore, if hence the multiplier  $L$  could be found in general, one would have the universal resolution of all differential equation of degree one; it is certainly not possible to hope for this. Therefore, we have to be content, if for the various cases, and many classes of differential equations, we can investigate factors of this kind. But there are two classes of equations, for which such factors can be found conveniently, of which the one comprehends the equations, in which the the one variable never raises higher than one dimension; the other class on the other hand is the one of homogeneous equations. But except for these two classes there are many other cases, in which the invention of such a factor can be done, to have examined which will not be useless, since this seems to open the only way to develop and expand that branch of Analysis, which is still desired. Therefore, I decided to collect many classes of equations, which can be rendered integrable by means of a multiplier of this kind.

## PROBLEM 2

§21 Having known one single multiplier  $L$ , which renders the formula  $Mdx + Ndy$  integrable, to find infinitely many other multipliers, which have the same use.

## SOLUTION

Because the formula  $L(Mdx + Ndy)$  by assumption is integrable, let its integral be  $= z$ , such that it is

$$dz = L(Mdx + Ndy),$$

where  $z$  is a certain function of  $x$  and  $y$ . Now, let  $Z$  denote any function of  $z$ , and since the formula  $Zdz$  is also integrable, because of

$$Zdz = LZ(Mdx + Ndy)$$

it is manifest that the propounded formula  $Mdx + Ndy$  also becomes integrable, if it is multiplied by  $LZ$ . Therefore, having found one multiplier  $L$ , which renders the formula  $Mdx + Ndy$  integrable, from it innumerable other factors  $LZ$  can be found, which will enable the same by taking any arbitrary function of the integral

$$\int L(Mdx + Ndy)$$

for  $Z$ .

#### COROLLARY 1

§22 Therefore, having propounded any differential equation  $Mdx + Ndy$ , not only one but even infinitely many multipliers are given which render it integrable. But it suffices to have found one of them, since all remaining ones are determined by this one.

#### COROLLARY 2

§23 Therefore, if one has the differential equation

$$Mdx + Ndy = 0,$$

it can be rendered integrable in infinitely many ways. But no matter whether one takes the multiplier  $L$ , or any other one  $LZ$ , the found integral equation reduces to the same; for, since that factor  $L$  yields  $z = \text{Const.}$ , but this on the other hand  $\int Zdz = \text{Const.}$ , they agree, since  $\int Zdz$  is a function of  $z$ .

#### EXAMPLE 1

§24 To find all multipliers, which render this formula

$$\alpha y dx + \beta x dy$$

integrable.

One multiplier solving this problem is obvious, namely  $\frac{1}{xy}$ . Therefore, let  $L = \frac{1}{xy}$ , and further

$$dz = \frac{\alpha y dx + \beta x dy}{xy} = \frac{\alpha dx}{x} + \frac{\beta dy}{y},$$

whence by integrating it arises

$$z = \alpha \log x + \beta \log y = \log x^\alpha y^\beta.$$

Now, let  $Z$  denote any function of  $z = \log x^\alpha y^\beta$ , this means of  $x^\alpha y^\beta$ , and all multipliers in question will be contained in this general form

$$\frac{1}{xy} \text{funct. } x^\alpha y^\beta.$$

Therefore, one will find simpler multipliers, if instead of the function any function of  $x^\alpha y^\beta$  is taken; and so the formula  $\alpha y dx + \beta x dy$  is rendered integrable by this further extending multiplier  $x^{\alpha n-1} y^{\beta n-1}$ . If more composite ones are desired, one will be able to combine any number of formulas of this kind that one has

$$A x^{\alpha n-1} y^{\beta n-1} + B x^{\alpha m-1} y^{\beta m-1} + \text{etc.}$$

### EXAMPLE 2

§25 To find all multipliers which renders this differential formula

$$\alpha x^{\mu-1} y^{\nu} dx + \beta x^{\mu} y^{\nu-1} dy$$

integrable.

Here, again one multiplier immediately reveals itself to us

$$L = \frac{1}{x^{\mu} y^{\nu}},$$

which yields

$$dz = \frac{\alpha dx}{x} + \frac{\beta dy}{y},$$

whence it is

$$z = \alpha \log x + \beta \log y = \log x^{\alpha} y^{\beta}.$$

Therefore, having put  $Z$  for any function of  $x^{\alpha} y^{\beta}$ , all multipliers will be contained in this general expression

$$\frac{Z}{x^{\mu} y^{\nu}} = \frac{1}{x^{\mu} y^{\nu}} \text{funct. } x^{\alpha} y^{\beta}.$$

If instead of this function one takes any arbitrary power of  $x^{\alpha n} y^{\beta n}$ , one will hence obtain innumerable multipliers, consisting of one single term  $x^{\alpha n - \mu} y^{\beta n - \nu}$ , by taking any arbitrary numbers for  $n$ .

### SCHOLIUM

§26 Therefore, it can happen that two or more differential formulas of this kind

$$\alpha x^{\mu-1} y^{\nu} dx + \beta x^{\mu} y^{\nu-1} dy$$

receive a common multiplier: If this happens, the differential equation composed of formulas of this kind as terms can be rendered integrable, if this common multiplier is applied. Let us expand this case already considered once.

### PROBLEM 3

§27 Let this differential equation be propounded:

$$\alpha y dx + \beta x dy + \gamma x^{\mu-1} y^{\nu} dx + \delta x^{\mu} y^{\nu-1} dy = 0,$$

whose integral is to be found.

### SOLUTION

To find an appropriate multiplier such that this equation is rendered integrable, consider both terms separately. And we certainly saw the first term  $\alpha y dx + \beta x dy$  to be rendered integrable by this multiplier

$$x^{\alpha n-1} y^{\beta n-1},$$

but the second term  $\gamma x^{\mu-1} y^{\nu} dx + \delta x^{\mu} y^{\nu-1} dy$  by this one

$$x^{\gamma m-\mu} y^{\delta m-\nu}.$$

Since now for  $n$  and  $m$  any arbitrary numbers can be taken, these two factors can be made equal; hence it is

$$\alpha n - 1 = \gamma m - \mu \quad \text{and} \quad \beta n - 1 = \delta m - \nu$$

and hence

$$n = \frac{\gamma m - \mu + 1}{\alpha} = \frac{\delta m - \nu + 1}{\beta},$$

and hence one obtains



$$m = \frac{\alpha\nu - \beta\mu - \alpha + \beta}{\alpha\delta - \beta\gamma} \quad \text{and} \quad n = \frac{\gamma\nu - \delta\mu - \gamma + \delta}{\alpha\delta - \beta\gamma}.$$

Having found these values for  $m$  and  $n$ , this common multiplier will give this integral equation:

$$\frac{1}{n}x^{\alpha n}y^{\beta n} + \frac{1}{m}x^{\gamma m}y^{\delta m} = \text{Const.}$$

#### COROLLARY 1

§28 Therefore, this integral equation is always algebraic, if for  $m$  and  $n$  the true values are found. Therefore, only the cases need a particular reduction, in which the number  $m$  and  $n$  go over into infinity or vanish.

#### COROLLARY 2

§29 But the two numbers  $m$  and  $n$  become infinite, if it was  $\alpha\delta = \beta\gamma$ . But in this case the differential is resolved into two factors, and acquires this form

$$(\alpha y dx + \beta x dy)\left(1 + \frac{\gamma}{\alpha}x^{\mu-1}y^{\nu-1}\right) = 0$$

and hence it will be

$$\text{either } \alpha y dx + \beta x dy = 0, \quad \text{or} \quad 1 + \frac{\gamma}{\alpha}x^{\mu-1}y^{\nu-1} = 0,$$

none of which two resolutions is of any difficulty.

#### COROLLARY 3

§30 But if it is  $n = 0$ , or

$$\gamma(\nu - 1) = \delta(\mu - 1),$$

consider the number  $n$  as very small, and because it is by means of a convergent series

$$x^{\alpha n} = 1 + \alpha n \log x + \frac{1}{2}\alpha^2 n^2 (\log x)^2 + \text{etc.} \quad \text{and} \quad y^{\beta n} = 1 + \beta n \log y + \frac{1}{2}\beta^2 n^2 (\log y)^2 + \text{etc.},$$

it will be

$$\frac{1}{n}x^{\alpha n}y^{\beta n} = \frac{1}{n} + \alpha \log x + \beta \log y = \log x^{\alpha}y^{\beta}$$

having involved the first part  $\frac{1}{n}$  into the constant. Therefore, in this case the integral equation will be:

$$\log x^{\alpha}y^{\beta} + \frac{1}{m}x^{\gamma m}y^{\delta m} = \text{Const.}$$

#### COROLLARY 4

§31 Therefore, for this case set

$$\mu = \gamma k + 1 \quad \text{and} \quad \nu = \delta k + 1,$$

that one has this differential equation:

$$\alpha y dx + \beta x dy + \gamma x^{\gamma k} \delta^{\delta k + 1} dx + \delta x^{\gamma k + 1} y^{\delta k} dy = 0,$$

and because it is

$$m = \frac{\alpha \delta k - \beta \gamma k}{\alpha \delta - \beta \gamma} = k,$$

the integral equation will be

$$\log x^{\alpha}y^{\beta} + \frac{1}{k}x^{\gamma k}y^{\delta k} = \text{Const.}$$

#### COROLLARY 5

§32 In similar manner, if it was  $m = 0$  or

$$\alpha(\nu - 1) = \beta(\mu - 1),$$

because of

$$\frac{1}{m}x^{\gamma m}y^{\delta m} = \log x^{\gamma}y^{\delta},$$

if one puts  $\mu = \alpha k + 1$  and  $\nu = \beta k + 1$ , whence it is

$$n = \frac{\gamma \beta k - \delta \alpha k}{\alpha \delta - \beta \gamma} = -k,$$

the integral of this equation

$$\alpha y dx + \beta x dy + \gamma x^{\alpha k} y^{\beta k + 1} dx + \delta x^{\alpha k + 1} y^{\beta k} dy = 0$$

will be

$$-\frac{1}{k} x^{-\alpha k} y^{-\beta k} + \log x^{\gamma} y^{\delta} = \text{Const.}$$

#### SCHOLIUM

§33 But a resolution of this kind in members, which are rendered integrable by means of the same multiplicator, does not extend to equations of all classes. For, it can certainly happen that the whole equation multiplied by a quantity becomes integrable, although no parts of it is integrable separately, from which one must not attribute too much to this treatment, which I did here.

#### PROBLEM 4

§34 If this differential equation is propounded

$$P dx + Q y dx + R dy = 0,$$

where  $P$ ,  $Q$  and  $R$  denote any arbitrary functions of  $x$ , such that the other variable  $y$  does not have more than one dimension, to find the multiplicator, which renders it integrable.

#### SOLUTION

Having compared this equation to the form  $M dx + N dy = 0$  it will be

$$M = P + Qy \quad \text{and} \quad N = R,$$

whence it will be

$$\left(\frac{dM}{dy}\right) = Q \quad \text{and} \quad \left(\frac{dN}{dx}\right) = \frac{dR}{dx}.$$

Now set  $L$  for the multiplicator in question, and let

$$dL = p dx + q dy,$$

and this equation must be satisfied

$$\frac{Np - Mq}{L} = Q - \frac{dR}{dx} = \frac{Rp - (P + Qy)q}{L}.$$

Because now  $Q - \frac{dR}{dx}$  is a function of  $x$  only, one will be able to take a function of  $x$  only for  $L$  also, such that it is  $q = 0$ , and  $dL = p dx$ ; hence it will be:

$$Q - \frac{dR}{dx} = \frac{Rp}{L}, \quad \text{or} \quad Q dx - dR = \frac{R dL}{L}$$

and hence

$$\frac{dL}{L} = \frac{Q dx}{R} - \frac{dR}{R}.$$

Hence by integrating one will have

$$\log L = \int \frac{Q dx}{R} - \log R,$$

and having taken  $e$  for the number whose hyperbolic logarithm is unity it arises

$$L = \frac{1}{R} e^{\int \frac{Q dx}{R}}.$$

But having found this multiplier the integral equation will be:

$$\int \frac{P dx}{R} e^{\int \frac{Q dx}{R}} + y e^{\int \frac{Q dx}{R}} = \text{Const.}$$

#### COROLLARY 1

§35 If the equation has the propounded form, it, before it is treated this way, can be divided by  $R$  that it obtains this form

$$P dx + Q y dx + dy = 0,$$

or one can immediately assume  $R = 1$ , having done which the multiplier will be  $e^{\int Q dx}$ , and the integral equation

$$\int e^{\int Q dx} P dx + e^{\int Q dx} y = \text{Const.}$$

## COROLLARY 2

§36 If one puts this integral

$$\int e^{\int Q dx} P dx + e^{\int Q dx} y = z,$$

such that  $z$  is a certain function of two variables, but then  $Z$  denotes any arbitrary function of  $z$ , all multipliers, which render the formula

$$P dx + Q dy + dy$$

integrable, are contained in this general form  $e^{\int Q dx}$ .

## PROBLEM 5

§37 If this differential equation is propounded:

$$P y^n dx + Q dy dx + R dy = 0,$$

where  $P$ ,  $Q$  and  $R$  denote any functions of  $x$ , to find a multiplier, which renders it integrable.

## SOLUTION

Therefore, it will be  $M = P y^n + Q y$  and  $N = R$ , and hence

$$\left( \frac{dM}{dy} \right) = n P y^{n-1} + Q \quad \text{and} \quad \left( \frac{dN}{dx} \right) = \frac{dR}{dx}.$$

Hence having put the multiplier in question  $L$  and

$$dL = p dx + q dy,$$

from the things found before it will be:

$$\frac{R p - P y^n q - Q y q}{L} = n P y^{n-1} + Q - \frac{dR}{dx}.$$

Assume  $L = S y^m$ , while  $S$  is a function of  $x$  only, then it will be

$$p = \frac{y^m dS}{dx} \quad \text{and} \quad q = m S y^{m-1},$$

having substituted which values, it will arise:

$$\frac{RdS}{Sdx} - mPy^{n-1} - mQ = nPy^{n-1} + Q - \frac{dR}{dx}.$$

That this equation can hold, one has to take  $m = -n$ , and it will be

$$\frac{RdS}{Sdx} = (1-n)Q - \frac{dR}{dx}, \quad \text{or} \quad \frac{dS}{S} = \frac{(1-n)Qdx}{R} - \frac{dR}{R}.$$

Hence, because by integration it arises

$$S = \frac{1}{R} e^{(1-n) \int \frac{Qdx}{R}},$$

because of  $m = -n$  the multiplier in question will be:

$$L = \frac{y^{-n}}{R} e^{(1-n) \int \frac{Qdx}{R}}$$

and the integral equation will be

$$\frac{y^{1-n}}{1-n} e^{(1-n) \int \frac{Qdx}{R}} + \int \frac{Pdx}{R} e^{(1-n) \int \frac{Qdx}{R}} = \text{Const.}$$

#### COROLLARY 1

§38 If it is  $n = 0$ , we have the case treated before of the equation

$$Pdx + Qdx + Qydx + Rdy = 0,$$

which by means of the multiplier

$$\frac{1}{R} e^{\int \frac{Qdx}{R}}$$

is rendered integrable; and the integral equation of it is

$$ye^{\int \frac{Qdx}{R}} + \int \frac{Pdx}{R} e^{\int \frac{Qdx}{R}} = \text{Const.}$$

#### COROLLARY 2

§39 But let  $n = 1$ , that the differential equation is:

$$Pydx + Qdydx + Rdy = 0$$

the multiplier because of  $1 - n = 0$  will be  $\frac{1}{Ry}$ ; that the equation is reduced to this form

$$\int \frac{(P + Q)dx}{R} + \log y = \text{Const.}$$

#### SCHOLIUM

§40 Additionally, this problem is easily deduced from the preceding problem. For, divide the propounded differential equation by  $y^n$ , and one will have:

$$Pdx + Qy^{1-n}dx + Ry^{-n}dy = 0.$$

Put  $y^{1-n} = z$ , it will be  $(1 - n)y^{-n}dy = dz$ , and so the equation goes over into this one:

$$Pdx + Qzdx + \frac{1}{1 - n}Rdz = 0,$$

which agrees to the equation of the preceding problem. Therefore, because these two equations are to be referenced to the case, in which the one variable never ascends higher than one dimension, we have completed it by this method of multipliers. Therefore, I proceed to another class, the class of homogeneous differential equations, which are known that they can also be treated by this method. But for this it is necessary to give a lemma, in which the nature of homogeneous functions is contained, in advance, if we want to derive the operation on first principles.

#### LEMMA

§41 If  $V$  was a homogeneous function, in which the two variables  $x$  and  $y$  constitute  $n$  dimensions everywhere, its differential

$$dV = Pdx + Qdy$$

will be of such a nature that it is

$$Px + Qy = nV.$$

### DEMONSTRATION

Put  $y = xz$ , and the function  $V$  will obtain a form of this kind  $x^n Z$ , where  $Z$  is a certain function of  $z$  only. Therefore, it will hence be

$$dV = nx^{n-1}Zdx + x^n dZ.$$

Reduce these two variables  $x$  and  $z$  also to the propounded differential  $dV = Pdx + Qdy$ , and because it is

$$dy = zdx + xdz,$$

it will be

$$dV = (P + Qz)dx + Qxdz;$$

therefore, it is necessary that it is

$$nx^{n-1}Z = P + Qz,$$

and by multiplying by  $x$  on both sides:

$$nx^n Z = nV = Px + Qxz = Px + Qy,$$

such that it is  $Px + Qy = nV$ .

### COROLLARY 1

§42 Therefore, since we have the two equations:

$$dV = Pdx + Qdy \quad \text{and} \quad nV = Px + Qy,$$

hence the two functions  $P$  and  $Q$  can be defined; for, one will find:

$$P = \frac{ydV - nVdy}{ydx - xdy} \quad \text{and} \quad Q = \frac{nVdx - xdV}{ydx - xdy}.$$

### COROLLARY 2

§43 Therefore, as often as  $V$  is a homogeneous function of  $n$  dimensions, so often because of



$$P = \left( \frac{dV}{dx} \right) \quad Q = \left( \frac{dV}{dy} \right)$$

it will be

$$\left( \frac{dV}{dx} \right) = \frac{y dV - nV dy}{y dx - x dy} \quad \text{and} \quad \left( \frac{dV}{dy} \right) = \frac{nV dx - x dV}{y dx - x dy},$$

where it is to be noted that in these fractions the differentials cancel each other, or both numerators will be divisible by  $y dx - x dy$ .

#### PROBLEM 6

§44 Having propounded the differential equation

$$M dx + N dy = 0,$$

in which  $M$  and  $N$  are homogeneous functions of  $x$  and  $y$ , both of the same number of dimensions, to find the multiplicator, which renders the equation integrable.

#### SOLUTION

Let  $n$  be the number of dimensions corresponding to both functions  $M$  and  $N$ , and by means of the preceding paragraph it will be

$$\left( \frac{dM}{dy} \right) = \frac{nM dx - x dM}{y dx - x dy} \quad \text{and} \quad \left( \frac{dN}{dx} \right) = \frac{y dN - nN dy}{y dx - x dy}$$

and hence

$$\left( \frac{dM}{dy} \right) - \left( \frac{dN}{dx} \right) = \frac{n(M dx + N dy) - x dM - y dN}{y dx - x dy}.$$

Now, it is easily concluded that a multiplicator is given which is also a homogeneous function of  $x$  and  $y$ . Therefore, let  $L$  be such a homogeneous function of  $m$  dimensions. Hence, if in § 19 one puts

$$dL = P dx + Q dy,$$

it will be [§ 42]

$$P = \frac{y dL - m L dy}{y dx - x dy} \quad \text{and} \quad Q = \frac{m L dx - x dL}{y dx - x dy}$$

and hence, because it must be according to § 19

$$\frac{NP - MQ}{L} = \left( \frac{dM}{dy} \right) - \left( \frac{dN}{dx} \right),$$

by multiplying by  $y dx - x dy$  on both side one will obtain:

$$\frac{Ny dL - m L N dy - m L M dx + M x dL}{L} = n(M dx + N dy) - x dM - y dN,$$

whence one finds:

$$\frac{dL}{L} = \frac{(m+n)(M dx + N dy) - x dM - y dN}{M x + N y},$$

which formula manifestly becomes integrable having put  $m+n = -1$ , having done which it will be

$$\log L = -\log(Mx + Ny).$$

Therefore, one will have the multiplier in question

$$L = \frac{1}{Mx + Ny}.$$

#### COROLLARY 1

**§45** Therefore, having propounded the homogeneous differential equation  $M dx + N dy = 0$ , is will most easily be reduced to integrability, since the formula

$$\frac{M dx + N dy}{M x + N y}$$

is integrable, whose integral, by means of the method given above, will give the integral equation is question.

## COROLLARY 2

§46 A inconvenience arises only in the case, where it is  $Mx + Ny = 0$ , as it happens in the equation  $ydx - xdy = 0$ , which would have to be divided by

$$xy - xy = 0 \cdot xy.$$

But since any multiple of this divisor equally satisfies, the divisor  $xy$  will also solves the task, as it is perspicuous per se.

## SCHOLIUM

§47 There is a very well known method, by which the most ingenious Joh. Bernoulli once taught to render all homogeneous differential equations separable. Having propounded an equation of this kind

$$Mdx + Ndy = 0$$

in which  $M$  and  $N$  shall be homogeneous functions of  $n$  dimension, he tells us to put  $y = ux$ , having done which the functions  $M$  and  $N$  will obtain forms of this kind, that it is

$$M = x^n U \quad \text{and} \quad N = x^n V,$$

while  $U$  and  $V$  are functions of  $u$  only. Therefore, the propounded equation, divided by  $x^n$  will go over into this one:

$$Udx + Vdy = 0.$$

But because it is  $dy = udx + xdu$ , we will have

$$Udx + Vudx + Vxdu = 0,$$

which divided by  $x(U + Vu)$  becomes separable, or this form

$$\frac{(U + Vu)dx + Vxdu}{x(U + Vu)}$$

integrable. But it is

$$(U + Vu)dx + Vxdu = \frac{1}{x^n}(Mdx + Ndy)$$

and

$$x^n(U + Vu) = M + Nu.$$

Therefore, this formula will be integrable:

$$\frac{Mdx + Ndy}{x(M + Nu)} = \frac{Mdx + Ndy}{Mx + Ny} \quad \text{because of } ux = y.$$

Therefore, having explained these two classes of equations, which can be rendered integrable by means of suitable multipliers, let us see, to which other classes the same method can be extended: And at first I observe that all differential equations which can be integrated by other methods can also be treated by this method by means of a suitable multiplier, which will be explained more clearly in the following problem.

#### PROBLEM 7

**§48** Having propounded the differential equation  $Mdx + Ndy = 0$ , if its complete integral equation was found, to assign all multipliers, which render the differential equation integrable.

#### SOLUTION

Since the complete integral equation involves an arbitrary constant  $C$ , which is not in the differential equation, now matter how intricate it might be, find its value by resolution of the equation, which shall be  $C = V$ , and  $V$  will be a function of  $x$  and  $y$  which additionally contains constants of the differential equation in it. Then differentiate this equation  $C = V$ , and so  $0 = dV$  will arise. And now it is necessary that  $dV$  has the propounded form itself as a divisor. Therefore, let it be

$$dV = L(Mdx + Ndy),$$

and  $L$  will be the suitable multiplier, which renders the propounded differential equation integrable. Further, since, while  $Z$  denotes any function of  $V$ , the formula

$$ZdV = LZ(Mdx + Ndy)$$

is also integrable, the expression  $LZ$  will include all multipliers, by which the propounded differential equation  $Mdx + Ndy = 0$  becomes integrable.

### COROLLARY 1

§49 Therefore, as often as the complete integral of the differential equation  $Mdx + Ndy = 0$  can be assigned, so often not only one but completely all multipliers can be defined by which the equation is rendered integrable.

### COROLLARY 2

§50 Therefore, because by other methods the complete integrals of many differential equations were found, hence the method treated up to now, which was still applied only to two classes of equations, can be amplified significantly.

### SCHOLIUM

§51 Nevertheless, if we do not want to descend to very special cases, the differential equations, whose complete integrals can be assigned, are reduced to a tiny number. And first the differential equations of first degree contained in this form

$$dx(\alpha + \beta x + \gamma y) + dy(\delta + \epsilon x + \zeta y) = 0,$$

since which are easily reduced to homogeneous ones, can also be treated by this method of multipliers. Furthermore, this form is remarkable

$$dy + Pydx + Qyydx = Rdx,$$

if one single satisfying value of which is known, from it the complete integral can be found, whence in these cases it will be possible to assign appropriate multipliers. Thirdly, also the cases of this equation deserve it to be considered

$$dy + yydx = ax^m dx,$$

called the Riccati equation after its discoverer, in which it can be reduced to separability. Finally, there are the cases of this equation

$$ydy + Pydx = Qdx,$$

since which are integrable, they are accommodated to the investigation of multipliers. Hence this will open a new way to find the equation from a given form of the multipliers, which by means of them become integrable,

whence it might be possible to gain increments for Analysis not to be hoped for.

### PROBLEM 8

§52 Having propounded the differential equation of first degree:

$$(\alpha + \beta x + \gamma y)dx + (\delta + \epsilon x + \zeta y)dy = 0,$$

to find the multipliers, which render it integrable.

### SOLUTION

Reduce this equation to homogeneity by putting:

$$x = t + f \quad \text{and} \quad y = u + g,$$

that it arises

$$(\alpha + \beta f + \gamma g + \beta \gamma u)dt + (\delta + \epsilon f + \zeta g + \epsilon t + \zeta u)du = 0$$

which having put

$$\alpha + \beta f + \gamma g = 0 \quad \text{and} \quad \delta + \epsilon f + \zeta g = 0,$$

whence the quantities  $f$  and  $g$  are determined, becomes homogeneous, of course,

$$(\beta t + \gamma u)dt + (\epsilon t + \zeta u)du = 0;$$

and hence by means of the multiplier

$$\frac{1}{\beta t t + (\gamma + \epsilon) t u + \zeta u u}$$

it is rendered integrable. Hence having found the letters  $f$  and  $g$  the propounded equation will become integrable, if it is divided by

$$\beta(x - f)^2 + (\gamma + \epsilon)(x - f)(y - g) + \zeta(y - g)^2,$$

or by

$$\beta x x + (\gamma + \epsilon) x y + \zeta y y - (2\beta f + \gamma g + \epsilon g) x - (2\zeta g + \gamma f + \epsilon f) y$$

$$+\beta ff + (\gamma + \varepsilon)fg + \zeta gg.$$

But because it is

$$f = \frac{\alpha\zeta - \gamma\delta}{\gamma\varepsilon - \beta\zeta} \quad \text{and} \quad g = \frac{\beta\delta - \alpha\varepsilon}{\gamma\varepsilon - \beta\zeta},$$

the divisor in question will arise as:

$$\begin{aligned} & \beta xx + (\gamma + \varepsilon)xy + \zeta yy + \frac{\alpha\gamma\delta - \alpha\alpha\zeta + \alpha + \delta\varepsilon - \beta\delta\delta}{\gamma\varepsilon - \beta\zeta} \\ & + \frac{-2\alpha\beta\zeta + \beta\gamma\delta - \beta\delta\varepsilon + \alpha\gamma\varepsilon + \alpha\varepsilon\varepsilon}{\gamma\varepsilon - \beta\zeta}x + \frac{-2\beta\delta\zeta + \alpha\varepsilon\zeta - \alpha\gamma\zeta + \gamma\delta\varepsilon + \gamma\gamma\delta}{\gamma\varepsilon - \beta\zeta}y. \end{aligned}$$

But having found one divisor or multiplicator, from it one will easily find all possible ones.

### COROLLARY 1

§53 Therefore, the form of the divisor by means of which the differential equation

$$(\alpha + \beta x + \gamma y)dx + (\delta + \varepsilon x + \zeta y)dy = 0$$

is rendered integrable, it is

$$\beta xx + (\gamma + \varepsilon)yx + \zeta yy + Ax + By + C,$$

where the constants  $A, B, C$  were defined above.

### COROLLARY 2

§54 Since the found divisor also satisfies, if it is multiplied by  $\gamma\varepsilon - \beta\zeta$ , it is plain that in the case, in which it is  $\beta\zeta = \gamma\varepsilon$ , the divisor will be

$$(\alpha\varepsilon\varepsilon - \beta\delta\varepsilon + \beta\gamma\delta - \alpha\beta\zeta)x + (\gamma\gamma\delta - \alpha\gamma\delta - \alpha\gamma\zeta + \alpha\varepsilon\zeta - \beta\delta\zeta)y + \alpha\gamma\delta - \alpha\alpha\zeta + \alpha\delta\varepsilon - \beta\delta\delta$$

which having put

$$\beta = mf, \quad \gamma = nf, \quad \varepsilon = mg, \quad \zeta = ng,$$

goes over into

$$m(\alpha g - \delta f)(mg - nf)x + n(\alpha g - \delta f)(mg - nf)y + (\alpha g - \delta f)(\delta m - \alpha n).$$

### COROLLARY 3

§55 Hence, if the propounded equation was of this kind:

$$[\alpha + f(mx + ny)]dx + [\delta + g(mx + ny)]dy = 0,$$

it will be rendered integrable, if it is divided by

$$(mg - nf)(mx + ny) + \delta m - \alpha n$$

or by

$$mx + ny + \frac{\delta m - \alpha n}{mg - nf}.$$

But if it was  $mg - nf = 0$ , the propounded equation itself is already integrable.

### PROBLEM 9

§56 Having propounded this differential equation:

$$dy + Pydx + Qydy + Rdx = 0,$$

where  $P$ ,  $Q$  and  $R$  are functions of  $x$  only, if it is known that this equation is satisfied  $y = v$ , where  $v$  is a function of  $x$ , to find multipliers, which render this equation integrable.

### SOLUTION

Since the value  $y = v$  satisfies the equation, it will be

$$dy + Pvdx + Qvdy + Rdx = 0;$$

therefore, if one puts  $y = v + \frac{1}{z}$ , one will have

$$-\frac{dz}{zz} + \frac{Pdx}{z} + \frac{2Qvdx}{z} + \frac{Qdx}{zz} = 0$$

or



$$dz - (P - 2Qv)zdx - Qdx = 0,$$

which is rendered integrable by means of the multiplier

$$e^{-\int(P+2Qv)dx}.$$

Therefore, this multiplier multiplied by  $z$  will satisfy the propounded equation. Therefore, because  $z = \frac{1}{y-v}$  is a multiplier rendering the propounded equation integrable, it will be:

$$\frac{1}{(y-v)^2} e^{-\int(P+2Qv)dx}.$$

For the sake of brevity let it be

$$e^{-\int(P+2Qv)dx} = S.$$

Since the integral of the equation

$$dz - (P + 2Qv)zdx - Qdx = 0$$

is

$$Sz - \int QSdx = \text{Const.},$$

all multipliers in question will be contained in this form:

$$\frac{S}{(y-v)^2} \text{funct.} \left( \frac{S}{y-v} - \int QSdx \right),$$

where by assumption  $v$  is a known function of  $x$ , and hence also  $S = e^{-\int(P+2Qv)dx}$ .

#### COROLLARY 1

§57 Therefore, the multiplier, which reveals itself at first, is

$$\frac{S}{(y-v)^2},$$

then also

$$\frac{S}{S(y-v) - (y-v)^2 \int QSdx}$$

will be a multiplier, even if which contains the integral formula  $\int QSdx$ , it can often become simpler than that one.

### COROLLARY 2

§58 For, if  $S$  is an exponential quantity, it can happen that  $\int QSdx$  takes a form of this kind  $ST$ , where  $T$  is an algebraic function, in which case the multiplier will be

$$\frac{1}{y-v - (y-v)^2 T} = \frac{1}{(y-v)(1 - Ty + Tv)}$$

and hence algebraic, which cannot happen in the first form.

### COROLLARY 3

§59 Because in these two cases the multiplier is a fraction, only whose denominator the variable  $y$  goes into, and there does not ascend higher than a square, innumerable other multipliers of this kind can be exhibited: For, let  $\int QSdx = V$ , and it will be possible to multiply the denominator of the fraction  $\frac{S}{(y-v)^2}$  by

$$A + B \left( \frac{S}{y-v} - V \right) + C \left( \frac{S}{y-v} \right)^2,$$

and so a more general form of the multiplier will be:

$$\frac{S}{A(y-v)^2 + BS(y-v) - BV(y-v)^2 + CSS - 2CSV(y-v) + CVV(y-v)^2}$$

or:

$$\frac{S}{(A - BV + CVV)y^2 - (2Av - BS - 2BVv + 2CSV + 2CVVv)y + Avv - BSv - BSv - BVvv + CSS + 2CSVv + CV^2v^2}$$

COROLLARY 4

§60 Therefore, if this formula

$$\frac{dy + Pydx + Qyydx + Rdx}{Lyy + My + N}$$

was integrable, the denominator must be of such a nature that it is

$$SL = A - BV + CVV, \quad SM = S(B - 2CV) - 2v(A - BV + CVV)$$

and

$$SN = CSS - Sv(B - 2CV) + vv(A - BV + CVV)$$

where

$$dv + Pvdx + Qvvdv + Rdx = 0, \quad S = e^{-\int (P+2Qv)dx}$$

and  $V = \int QSdx$ .

PROBLEM 10

§61 Having propounded the preceding differential equation:

$$dy + Pydx + Qyydx + Rdx = 0$$

to find function  $L$ ,  $M$  and  $N$  of  $x$  that it divided by the formula

$$Lyy + My + N$$

becomes integrable.

SOLUTION

Therefore, because this formula must be integrable:

$$\frac{dy + dx(Py + Qyy + R)}{Lyy + My + N},$$

it is necessary by means of the general property, after we multiplied by

$$(Lyy + My + N)^2$$

that it is:

$$-\frac{yydL}{dx} - \frac{ydM}{dx} - \frac{dN}{dx} = \begin{matrix} +QMyy - 2RLy + NP \\ -PLyy + 2QNy - RM \end{matrix}$$

Hence for the determination of the functions  $L$ ,  $M$  and  $N$  we obtain these equations:

$$\begin{aligned} I. \quad dL &= PLdx - QMdx \\ II. \quad dM &= 2RLdx - 2QNdx \\ III. \quad dN &= RMdx - PNdx, \end{aligned}$$

from the first of which we deduce:

$$M = \frac{PL}{Q} - \frac{dL}{Qdx}$$

and from the second:

$$N = \frac{RL}{Q} - \frac{dM}{2Qdx'}$$

which values substituted for  $M$  and  $N$  in the third give:

$$dN = \frac{PdM}{2Q} - \frac{RdL}{Q}.$$

But because, having assumed the differential  $dx$  to be constant, it is

$$dM = \frac{PdL + LdP}{Q} - \frac{PLdQ}{QQ} - \frac{ddL}{Qdx} + \frac{dQdL}{QQdx'}$$

it will be

$$N = \frac{RL}{Q} - \frac{PdL}{2QQdx} - \frac{LdP}{2QQdx} + \frac{PLdQ}{2Q^3dx} + \frac{ddL}{2QQdx^2} - \frac{dQdL}{2Q^3dx^2}$$

and

$$dN = \frac{PPdL}{2QQ} + \frac{PLdP}{2QQ} - \frac{PPLdQ}{2Q^3} - \frac{PddL}{2QQdx} + \frac{PdQdL}{2Q^3dx} - \frac{RdL}{Q},$$

which therefore must be equal to the differential of the latter, whence it is

$$\begin{aligned}
0 = & QQd^3L - 3QdQddL - PPQQdLdx^2 - 2QQdPdLdx \\
& + 3dQ^2dL + 2PQdQdLdx - QdLddQ + 4Q^3RdLdx^2 \\
& - PQQLdPdx^2 + PPQLdQdx^2 - QQLdxddP + PQLdxddQ \\
& + 3QLdPdQdx - 3PLdQ^2dx + 2Q^3LdRdx^2 - 2Q^2RLdQdx^2.
\end{aligned}$$

But if this equation is multiplied by  $\frac{L}{Q^4}$ , it can be integrated, and its integral will be

$$\text{Const.} = \frac{LddL}{QQ} - \frac{LdLdQ}{Q^3} - \frac{dL^2}{2QQ} - \frac{PPLLdx^2}{2QQ} - \frac{LLdPdx}{QQ} + \frac{PLLdQdx}{Q^3} + \frac{2RLLdx^2}{Q},$$

which goes over into this form:

$$\begin{aligned}
2EQ^3dx^2 = & 2QLddL - 2LLdQ - QdL^2 - PPQLLdx^2 - 2QLLdPdx \\
& + 2PLLdQdx + 4QQRLLdx^2.
\end{aligned}$$

If one puts  $L = zz$ , the equation will obtain this form:

$$\frac{2EQ^3dx^2}{z^3} = 4Qddz - 4dQdz - z(PPQdx^2 + 2QdPdx - 2PdQdx - 4QQRdx^2).$$

#### COROLLARY 1

**§62** Therefore, as often as by means of the preceding problem the value of  $L$  can be assigned, so often the differential equation of third order found here and the one of second order, to which I reduced the latter, can be solved in general: This resolution, because it would be most difficult otherwise, is to be carefully noted.

#### COROLLARY 2

**§63** If  $v$  was a function of  $x$  of such a kind, which put instead of  $y$ , satisfies the equation

$$dy + Pydx + Qyydx + Rdx = 0,$$

take

$$S = e^{-\int(P+2Qv)dx}$$

and set  $V = \int QSdx$ , having done which for our differential equation of third order it will be

$$L = \frac{A - BV + CVV}{S},$$

since which value contains three arbitrary constants, it will therefore be the complete integral of that equation.

### COROLLARY 3

§63 If it is  $P = 0$ ,  $Q = 1$  and  $R$  any arbitrary function of  $x$ , the differential equation of third degree will obtain the form:

$$0 = d^3L + 4RdLdx^2 + 2LdRdx^2,$$

for the invention of whose complete integral at first find a function of  $x$  which shall be  $= v$  and which satisfies this equation

$$dy + vvdxdx + Rdx = 0;$$

then put

$$V = \int e^{-2\int vdx} dx,$$

and it will be

$$L = (A - BV + CVV)e^{+2\int vdx}.$$

### COROLLARY 4

§64 Therefore, the same integral will satisfy this differential equation of second degree:

$$2Edx^2 = 2LddL - dL^2 + 4RLLdx^2$$

and, having out  $L = zz$ , also this one:

$$\frac{Edx^2}{2z^3} = ddz + Rzdx^2,$$

for which it therefore is

$$z = e^{+\int v dx} \sqrt{A - BV + CVV}.$$

#### SCHOLIUM

§65 Therefore, this integration, which can hardly be achieved from other principles, completely deserves some consideration. Hence we obtain the complete integration of the following sufficiently far-extending differential equation second order:

$$ddz + Sdx dz + Tz dx^2 = \frac{Edx^2}{z^3} e^{-2\int S dx}.$$

For, first find the value of  $v$  from this differential equation of first degree;

$$dv + v dx + S dx + T dx = 0,$$

having found which therefore for the sake of brevity put

$$V = \int e^{-2\int v dx - \int S dx} dx$$

and it will be

$$z = e^{\int v dx} \sqrt{A + BV + CVV},$$

if only the arbitrary constants  $A, B, C$  are taken in such a way that it is

$$AC - \frac{1}{4}BB = E,$$

and so still two constant are still arbitrary, as the nature of a complete integration requires it.

#### EXAMPLE 1

§66 Let this differential equation be propounded

$$dy + y dx + yy dx - \frac{dx}{x} = 0,$$

whose multipliers, which render it integrable, are to be investigated.

Therefore, by transferring Problem 9 to this, it will be

$$P = 1, \quad Q = 1, \quad \text{and} \quad R = -\frac{1}{x},$$

and since the value  $y = \frac{1}{x}$  satisfies this equation, it will be  $v = \frac{1}{x}$ . Hence it will be

$$S = e^{-\int(1+\frac{2}{x})dx} = \frac{1}{xx}e^{-x}$$

and one will have the multiplier, which reveals itself at first,

$$= e^{-x} \frac{1}{(xy - 1)^2}.$$

But it is further possible to multiply this by any arbitrary function of this form

$$e^{-x} \frac{1}{x(xy - 1)} - \int e^{-x} \frac{dx}{xx};$$

but because this formula cannot be integrated, no other suitable multipliers can be assigned. Therefore, because of the first this formula is integrable:

$$e^{-x} \frac{1}{(xy - 1)^2} \left( dy + ydx + yydx - \frac{dx}{x} \right),$$

whose, if  $x$  is assumed to be constant, integral is

$$\frac{-e^x}{x(xy - 1)} + X,$$

which differentiated, having put  $y$  to be constant, yields

$$\frac{e^{-x}dx(xxy + 2xy - x - 1)}{xx(xy - 1)^2} + dX,$$

which must become equal to the other term

$$\frac{e^{-x}}{(xy - 1)^2} \left( ydx + yydx - \frac{dx}{x} \right),$$

whence it is

$$dX = \frac{e^{-x}dx}{xx(xy - 1)^2} (xxyy - 2xy + 1) = e^{-x} \frac{dx}{xx};$$

and so the complete integral of our equation is



$$\frac{-e^{-x}}{x(xy-1)} + \int e^{-x} \frac{dx}{xx} = \text{Const.}$$

EXAMPLE 2

§67 To find suitable multipliers, which render this equation integrable:

$$dy + yydx - \frac{adx}{(\alpha + \beta x + \gamma xx)^2} = 0.$$

A singular case satisfying this equation is

$$y = \frac{k + \gamma x}{\alpha + \beta x + \gamma xx} = v$$

while

$$k = \frac{1}{2}\beta \pm \sqrt{\frac{1}{4}\beta\beta - \alpha\gamma + a}.$$

Because now it is  $P = 0$  and  $Q = 1$ , it will be

$$S = e^{-\int \frac{2kdx + 2\gamma xdx}{\alpha + \beta x + \gamma xx}}$$

or having put for the sake of brevity

$$\pm \sqrt{\frac{1}{4}\beta\beta - \alpha\gamma + a} = \frac{1}{2}n$$

it will be

$$S = \frac{1}{\alpha + \beta x + \gamma xx} e^{-\int \frac{ndx}{\alpha + \beta x + \gamma xx}}$$

and

$$\int Sdx = -\frac{1}{n} e^{-\int \frac{ndx}{\alpha + \beta x + \gamma xx}}.$$

Therefore, the multiplier found first is

$$e^{-\int \frac{ndx}{\alpha + \beta x + \gamma xx}} \cdot \frac{\alpha + \beta x + \gamma xx}{((\alpha + \beta x + \gamma xx)y - k - \gamma x)^2},$$

which can further be multiplied by any arbitrary function of this kind

$$e^{-\int \frac{ndx}{\alpha + \beta x + \gamma xx}} \left( \frac{1}{(\alpha + \beta x + \gamma xx)y - k - \gamma x} + \frac{1}{n} \right).$$

Therefore, multiply it by

$$e^{\int \frac{ndx}{\alpha + \beta x + \gamma xx}} \cdot \frac{(\alpha + \beta x + \gamma xx)y - k - \gamma x}{(\alpha + \beta x + \gamma xx)y + n - k - \gamma x}$$

and this algebraic multiplicator will arise:

$$\frac{\alpha + \beta x + \gamma xx}{((\alpha + \beta x + \gamma xx)y - k - \gamma x)((\alpha + \beta x + \gamma xx)y + n - k - \gamma x)},$$

which is reduced to this form:

$$\frac{1}{(\alpha + \beta x + \gamma xx) \left( y - \frac{2\gamma x + \beta + \sqrt{\beta\beta - 4\alpha\gamma + 4a}}{2(\alpha + \beta x + \gamma xx)} \right) \left( y + \frac{2\gamma x + \beta + \sqrt{\beta\beta - 4\alpha\gamma + 4a}}{2(\alpha + \beta x + \gamma xx)} \right)}.$$

But the complete integral of the equation is

$$e^{-\int \frac{ndx}{\alpha + \beta x + \gamma xx}} \frac{(\alpha + \beta x + \gamma xx)y + n - k - \gamma x}{(\alpha + \beta x + \gamma xx)y - k - \gamma x} = \text{Const.},$$

while  $n = \sqrt{\beta\beta - 4\alpha\gamma + 4a}$  and  $k = \frac{\beta + n}{2}$ .

Hence the complete integral equation will be

$$e^{-\int \frac{ndx}{\alpha + \beta x + \gamma xx}} \cdot \frac{2(\alpha + \beta x + \gamma xx)y + n - \beta - 2\gamma x}{2(\alpha + \beta x + \gamma xx)y - n - \beta - 2\gamma x} = \text{Const.},$$

whose nature is manifest, as long as

$$n = \sqrt{\beta\beta - 4\alpha\gamma + 4a}$$

is a real number.

But if the value of  $n$  is imaginary, say  $n = m\sqrt{-1}$ , because of

$$e^{p\sqrt{-1}} = \cos p + \sqrt{-1} \sin p,$$

the integral can be reduced to reality this way. Let

$$-\int \frac{dx}{\alpha + \beta x + \gamma x^2} = p \quad \text{and} \quad 2(\alpha + \beta x + \gamma x^2)y - \beta - 2\gamma x = q,$$

and it will be:

$$(\cos p + \sqrt{-1} \sin p) \cdot \frac{q + m\sqrt{-1}}{q - m\sqrt{-1}} = \text{Const.} = A + B\sqrt{-1},$$

hence it is:

$$q \cos p - m \sin p + (m \cos p + q \sin p)\sqrt{-1} = Aq + Bm + (Bq - Am)\sqrt{-1},$$

equate the real and imaginary terms separately:

$$q \cos p - m \sin p = Aq + Bm, \quad m \cos p + q \sin p = Bq - Am,$$

which two equations agree, if one takes  $A^2 + B^2 = 1$ . Therefore, let the arbitrary constant be  $A = \cos \theta$  that it is  $B = \sin \theta$  and in the case in which it is  $\sqrt{\beta^2 - 4\alpha\gamma} = m\sqrt{-1}$ , the real equation will be

$$q \cos p - m \sin p = q \cos \theta + m \sin \theta \quad \text{or} \quad q = \frac{m(\sin p + \sin \theta)}{\cos p - \cos \theta} = m \cot \frac{\theta - p}{2}.$$

Hence the complete integral equation of the differential equation

$$dy + ydx + \frac{(m^2 + \beta^2 - 4\alpha\gamma)dx}{4(\alpha + \beta x + \gamma x^2)^2} = 0,$$

having put

$$p = \int \frac{-m dx}{\alpha + \beta x + \gamma x^2},$$

is

$$2(\alpha + \beta x + \gamma x^2)y = \beta + 2\gamma x + m \cot \frac{\theta - p}{2}$$

or

$$y = \frac{\frac{1}{2}\beta + \gamma x + \frac{1}{2}m \cot \frac{\theta - p}{2}}{\alpha + \beta x + \gamma x^2},$$

or let  $\theta = 180^\circ - \zeta$ , and one will have

$$y = \frac{\frac{1}{2}\beta + \gamma x + \frac{1}{2}m \tan \frac{\zeta+p}{2}}{\alpha + \beta x + \gamma x^2}.$$

But in this case it is to be noted that the special integral from which we deduced all this becomes imaginary, what is nevertheless no obstruction that hence the complete integral can be exhibited in a real form.

### EXAMPLE 3

§68 *Having propounded the Riccati equation*

$$dy + ydx - ax^m dx = 0,$$

to find suitable multipliers for the cases of the exponent  $m$ , in which it can be separated.

Let  $y = v$  the satisfying value, and because it is

$$P = 0, \quad Q = 1 \quad \text{and} \quad R = -ax^m,$$

the first multiplier rendering the equation integrable will be

$$e^{-2 \int v dx} \frac{1}{(y - v)^2},$$

if it is multiplied by which, the complete integral becomes

$$e^{-2 \int v dx} \frac{1}{y - v} - \int e^{-2 \int v dx} dx = \text{Const.}$$

Hence, if  $Z$  denotes any arbitrary function of this quantity, all multipliers will be contained in this form:

$$e^{-2 \int v dx} \frac{Z}{(y - v)^2}.$$

Hence, if one puts

$$\int e^{-2 \int v dx} dx = V,$$

all multipliers contained in this form

$$\frac{1}{Ly + My + N}$$

will be obtained [§ 60], if one takes:

$$\begin{aligned} L &= e^{2 \int v dx} (A - BV + CVV) \\ M &= B - 2CV - 2ve^{2 \int v dx} (A - BV + CVV) \\ N &= Ce^{-2 \int v dx} - v(B - 2CV) + vve^{2 \int v dx} (A - BV + CVV). \end{aligned}$$

But this value of  $L$  at the same time is the complete integral of this differential equation of degree three:

$$0 = d^3L - 4ax^m dL dx^2 - 2maLx^{m-1} dx^3$$

and hence also of this one of degree two:

$$Edx^2 = 2LddL - dL^2 - 4aLLx^m dx^2$$

while

$$E = 4AC - BB.$$

#### SCHOLIUM

§69 Having studied the subject with more attention I even resolved the differential equation of third order by a direct method and detected that the same complete integral of it, which was assigned here, can be found. For, let this equation be propounded:

$$d^3L + 4RdL dx^2 + 2LdR dx^2 = 0,$$

where  $R$  is an arbitrary function of  $x$ , haven taken the differential  $dx$  to be constant. Now, I ask for a function of  $x$ , multiplied by which this differential equation becomes integrable. Let  $S$  be this function, and the integral of the equation

$$Sd^3L + 4SRdL dx^2 + 2SLdR dx^2 = 0$$

will be

$$SddL - dSdL + L(ddS + 4SRdx^2) = 2Cdx^2,$$

as long as it is

$$d^3S + 2SdRdx^2 + 4RdSdx^2 = 0.$$

It suffices, of course, to have taken a certain particularly satisfying value. But this equation multiplied by  $S$  having neglected the constant gives the integral:

$$SddS - \frac{1}{2}dS^2 + 2SSRdx^2 = 0.$$

Put  $S = e^{2 \int v dx}$ , and it will be

$$2dv + 2vvdx + 2Rdx = 0,$$

whence the task reduces to this that for  $v$  at least a particular value is investigated, which satisfies this differential equation of first degree:

$$dv + vvdx + Rdx = 0,$$

which I therefore assume as conceded. Hence our equation integrated one, because of  $S = e^{2 \int v dx}$  will be

$$ddL - 2vdx dL + L(2dvdx + 4vvdx^2 + 4Rdx^2) = 2Ce^{-2 \int v dx} dx^2.$$

Therefore, since because of

$$Rdx = -dv - vvdx$$

we have

$$ddL - 2vdx dL - 2Ldx dv = 2Ce^{-2 \int v dx} dx^2,$$

its integral manifestly is:

$$dL - 2Lvdx = Bdx + 2Cdx \int e^{-2 \int v dx} dx$$

and by multiplying the integral again by  $e^{-2 \int v dx}$  it will arise

$$e^{-2 \int v dx} L = A + B \int e^{-2 \int v dx} dx + 2C \int e^{-2 \int v dx} dx \int e^{-2 \int v dx} dx.$$

Hence, if for the sake of brevity one puts  $\int e^{-2\int v dx} dx = V$ , we will have

$$L = e^{2\int v dx} (A + BV + 2CVV)$$

completely as we found before.

#### PROBLEM 11

§70 Having propounded the Riccati equation

$$dy + yy dx = ax^m dx$$

to find its particular integrals in the cases in which it is separable.

#### SOLUTION

By putting  $a = cc$  and  $m = -4n$ , attribute this form to the equation:

$$dy + yy dx - ccx^{-4n} dx = 0.$$

For, since the question is about particular integrals, it is not important, whether they are real or not. But to find these cases, in which  $y$  can be expressed by means of a function of  $x$ , in an easier way and in one operation let us set

$$y = cx^{-2n} + \frac{dz}{z dx}$$

and having assumed  $dx$  to be constant, we will obtain this differential equation of second degree:

$$-2ncx^{-2n-1} dx + \frac{ddz}{z dx} + \frac{2cx^{-2n} dz}{z} = 0,$$

or

$$\frac{ddz}{dx^2} + \frac{2cdz}{x^{2n} dx} - \frac{2ncz}{x^{2n+1}} = 0,$$

whose value shall be assumed to be:

$$z = Ax^n + Bx^{3n-1} + Cx^{5n-2} + Dx^{7n-3} + Ex^{9n-4} + \text{etc.},$$

after having substituted which in the correct way we will obtain:

$$\begin{aligned}
0 = & n(n-1)Ax^{n-2} + (3n-1)(3n-2)Bx^{3n-3} + (5n-2)(5n-3)Cx^{5n-4} + \text{etc.} \\
& + 2ncAx^{-n-1} + 2(3n-1)cB \quad + \quad 2(5n-2)cC \quad + \quad 2(7n-3)cD \\
& - 2ncA \quad - \quad 2ncB \quad \quad \quad 2ncC \quad - \quad 2ncD
\end{aligned}$$

whence the assumed coefficients are determined this way:

$$\begin{aligned}
2(2n-1)cB + n(n-1)A = 0, & \quad B = \frac{-n(n-1)A}{2(2n-1)c} \\
2(4n-1)cC + (3n-1)(3n-2)B = 0, & \quad C = \frac{-(3n-1)(3n-2)B}{4(2n-1)c} \\
2(6n-3)cD + (5n-2)(5n-3)C = 0, & \quad D = \frac{-(5n-2)(5n-3)C}{6(2n-1)} \\
& \text{etc.}
\end{aligned}$$

Therefore, if one coefficient vanishes, all following ones at the same time vanish, what happens in these cases:

$$\begin{aligned}
n = 0, \quad n = \frac{1}{3}, \quad n = \frac{2}{5}, \quad n = \frac{3}{7}, \quad \text{etc.} \\
n = 1, \quad n = \frac{2}{3}, \quad n = \frac{3}{5}, \quad n = \frac{4}{7}, \quad \text{etc.}
\end{aligned}$$

Therefore, while  $i$  denotes any integer, as often as it was

$$n = \frac{i}{2i \pm i'}$$

so often the resolution of the equation can be exhibited. For, it will be

$$y = c^{-2n} + \frac{dz}{zdx},$$

where

$$z = Ax^n + Bx^{2n-1} + Cx^{5n-2} + Dx^{7n-3} + Ex^{9n-4} + \text{etc.}$$



Therefore, this particular value of  $y$  will arise:

$$y = cx^{-2n} + \frac{nAx^{n-1} + (3n-1)Bx^{3n-1} + (5n-2)Cx^{5n-3} + \text{etc.}}{Ax^n + Bx^{3n-1} + Cx^{5n-2} + \text{etc.}}$$

#### COROLLARY 1

§71 Therefore, if this particular value of  $y$  is called  $= v$ , a suitable multiplier of the propounded equation will be

$$= e^{-2 \int v dx} \cdot \frac{1}{(y-v)^2}.$$

And if one puts

$$\int e^{-2 \int v dx} dx = V,$$

having taken  $A = 0$  and  $C = 0$ , another simpler factor will be [§ 68]

$$\frac{1}{e^{2 \int v dx} V y y - (1 + 2ce^{2 \int v dx} V) y + v + vve^{2 \int v dx} V}.$$

#### COROLLARY 2

§72 But it is

$$\int v dx = \frac{-c}{(2n-1)x^{2n-1}} + \log(Ax^n + Bx^{2n-1} + Cx^{5n-1} + \text{etc.}),$$

whence it is

$$e^{-2 \int v dx} = e^{\frac{2c}{(2n-1)} x^{2n-1}} \frac{1}{(Ax^n + Bx^{3n-1} + Cx^{5n-2} + \text{etc.})^2},$$

from which further one finds the value of

$$V = \int e^{-2 \int v dx} dx,$$

if which was of this kind

$$e^{-2 \int v dx} T,$$

while  $T$  is an algebraic function, the superior multiplier will algebraic.

COROLLARY 3

§73 Having found the value  $v$ , or a particular integral of the propounded equation, hence one will immediately have the complete integral of the same, which will be:

$$\frac{e^{-2 \int v dx}}{y - v} - \int e^{-2 \int v dx} dx = \text{Const.}$$

CASE 1 IN WHICH IT IS  $n = 0$

§74 Therefore, for this equation

$$dy + yydx = ccdx,$$

because of  $B = 0, C = 0$  etc. a particular value will be  $y = c$ . Hence having put  $v = c$ , it will be

$$e^{-2 \int v dx} = e^{-2cx} \quad \text{and} \quad V = \int e^{-2 \int v dx} dx = -\frac{1}{2c} e^{-2cx};$$

hence the complete integral is

$$\frac{e^{-2cx}}{y - c} + \frac{y}{2c} e^{-2cx} = \text{Const..}$$

or

$$\frac{e^{-2cx(y+c)}}{y - c} = \text{Const.}$$

Further, because of

$$e^{2 \int v dx} V = -\frac{1}{2c} \quad \text{and} \quad v = c,$$

an algebraic multiplier will be:

$$\frac{1}{-\frac{1}{2c}yy + \frac{1}{2}c'}$$

which is reduced to

$$\frac{1}{yy - cc}$$

as it is perspicuous per se.

CASE 2 IN WHICH IT IS  $n = 1$

§75 Therefore, for this equation

$$dy + yydx = \frac{ccdxx}{x^4}$$

because of  $B = 0, C = 0$  a particular value will be

$$y = \frac{c}{xx} + \frac{1}{x}.$$

Hence having put

$$v = \frac{c}{xx} + \frac{1}{x},$$

it will be

$$e^{-2 \int v dx} = \frac{e^{\frac{2c}{x}}}{xx} \quad \text{and} \quad V = -\frac{1}{2c} e^{\frac{2c}{x}}.$$

Hence the complete integral is

$$\frac{e^{\frac{2c}{x}}}{xxy - x - c} + \frac{e^{\frac{2c}{x}}}{2c} = \text{Const.}$$

or

$$e^{\frac{2c}{x}} \cdot \frac{xxy - x + c}{xxy - x - c} = \text{Const.}$$

Further, because of

$$e^{2 \int v dx} V = -\frac{xx}{2c} \quad \text{and} \quad v = \frac{x+c}{xx},$$

one will have the algebraic multiplicator:

$$\frac{1}{xxyy - 2xy + 1 - \frac{cc}{xx}} = \frac{1}{(xy - 1)^2 - \frac{cc}{xx}}$$

or the propounded equation

$$dy + yydx - \frac{ccdxx}{x^4} = 0$$

becomes integrable, if it is divided by

$$(xy - 1)^2 - \frac{cc}{xx}.$$

CASE 3 IN WHICH IT IS  $n = \frac{1}{3}$

§76 Therefore, for this equation

$$dy + yydx - ccx^{-\frac{4}{3}}dx = 0$$

it is  $B = -\frac{A}{3c}$ ,  $C = 0$ , etc. whence a particular integral is

$$y = cx^{-\frac{2}{3}} + \frac{cx^{-\frac{2}{3}}}{3cx^{-\frac{1}{3}} - 1} = \frac{3ccx^{-\frac{1}{3}}}{3cx^{\frac{1}{3}} - 1} = v$$

and

$$e^{-2 \int v dx} = e^{-6cx^{\frac{1}{3}}} \frac{\text{Const.}}{\left(x^{\frac{1}{3}} - \frac{1}{3c}\right)^2} = e^{-6cx^{\frac{1}{3}}} \frac{1}{\left(3cx^{\frac{1}{3}} - 1\right)^2}$$

and hence

$$V = \int e^{-6cx^{\frac{1}{3}}} \frac{dx}{\left(3cx^{\frac{1}{3}} - 1\right)^2} = -e^{-6cx^{\frac{1}{3}}} \frac{3cx^{\frac{1}{3}} + 1}{18c^3 \left(3cx^{\frac{1}{3}} - 1\right)}.$$

Hence the complete integral is

$$\frac{e^{-6cx^{\frac{1}{3}}}}{\left(3cx^{\frac{1}{3}} - 1\right)^2 y - 3ccx^{-\frac{1}{3}} \left(3cx^{\frac{1}{3}} - 1\right)} + \frac{e^{-6cx^{\frac{1}{3}}} \left(3cx^{\frac{1}{3}} + 1\right)}{18c^4 \left(3cx^{\frac{1}{3}} - 1\right)} = \text{Const.}$$

or

$$e^{-6cx^{\frac{1}{3}}} \frac{y \left(1 + 3cx^{\frac{1}{3}}\right) + 3ccx^{-\frac{1}{3}}}{y \left(1 - 3cx^{\frac{1}{3}}\right) - 3ccx^{-\frac{1}{3}}} = \text{Const.}$$

Then, because of

$$e^{2 \int v dx} V = \frac{1 - 9ccx^{\frac{2}{3}}}{18c^3},$$

this divisor rendering the equation integrable will arise:

$$\left(y + 3ccx^{-\frac{1}{3}}\right)^2 - 9ccx^{\frac{2}{3}}yy.$$

CASE 4 IN WHICH IT IS  $n = \frac{2}{3}$

§77 Therefore, for this equation

$$dy + yydx - ccx^{-\frac{8}{3}}dx = 0$$

it is  $B = +\frac{A}{3c}$ ,  $C = 0$  etc., whence the particular integral is:

$$y = cx^{-\frac{4}{3}} \frac{2cx^{-\frac{1}{3}} + 1}{3cx^{\frac{2}{3}} + x} = \frac{3ccx^{-\frac{2}{3}} + 3cx^{-\frac{1}{3}} + 1}{3cx^{\frac{2}{3}} + x} = v$$

and

$$e^{-2 \int v dx} = e^{6cx^{-\frac{1}{3}}} \cdot \frac{1}{(3cx^{\frac{2}{3}} + x)^2};$$

hence it is further found:

$$V = \int \frac{e^{6cx^{-\frac{1}{3}}} dx}{(3cx^{\frac{2}{3}} + x)^2} = \frac{-e^{6cx^{-\frac{1}{3}}}(3cx^{\frac{2}{3}} - x)}{18c^3(3cx^{\frac{2}{3}} + x)}.$$

Hence the complete integral will be:

$$e^{6cx^{-\frac{1}{3}}} \frac{(x - 3cx^{\frac{2}{3}})y - 1 + 3cx^{-\frac{1}{3}} - 3ccx^{-\frac{2}{3}}}{(x + 3cx^{\frac{2}{3}})y - 1 - 3cx^{-\frac{1}{3}} - 3ccx^{-\frac{2}{3}}} = \text{Const.}$$

Then because of

$$e^{2 \int v dx} V = \frac{xx - 9ccx^{\frac{4}{3}}}{18c^3}$$

an algebraic divisor rendering the propounded equation integrable arises as:

$$((x + 3cx^{\frac{2}{3}})y - 1 - 3cx^{-\frac{1}{3}} - 3ccx^{-\frac{2}{3}})((x + 3cx^{\frac{2}{3}})y - 1 + 3cx^{-\frac{1}{3}} - 3ccx^{-\frac{2}{3}}).$$

CASE 5 IN WHICH IT IS  $n = \frac{2}{5}$

§78 Therefore, for the equation

$$dy + yydx - ccx^{-\frac{8}{5}} = 0$$

it will be

$$B = -\frac{3A}{5c}; \quad C = -\frac{B}{5c} = +\frac{3A}{35cc}; \quad D = 0 \quad \text{etc.}$$

and hence a particular integral:

$$y = cx^{-\frac{4}{5}} + \frac{\frac{2}{5}x^{-\frac{3}{5}} - \frac{1}{5} \cdot \frac{3}{5c}x^{-\frac{4}{5}}}{x^{\frac{2}{5}} - \frac{3}{5c}x^{\frac{1}{5}} + \frac{3}{25cc}} = cx^{-\frac{4}{5}} + \frac{10ccx^{-\frac{3}{4}} - 3cx^{-\frac{4}{5}}}{25ccx^{\frac{2}{5}} - 15cx^{\frac{1}{5}} + 3}$$

or

$$y = \frac{25c^3x^{-\frac{2}{5}} - 5ccx^{-\frac{3}{5}}}{25ccx^{\frac{2}{5}} - 15cx^{\frac{1}{5}} + 3} = v.$$

Hence the complete integral arises:

$$e^{-10cx^{\frac{1}{3}}} \cdot \frac{(3 + 15cx^{\frac{1}{5}} + 25ccx^{\frac{2}{5}})y + 5ccx^{-\frac{3}{5}} + 25c^3x^{-\frac{2}{5}}}{(3 - 15cx^{\frac{1}{5}} + 25ccx^{\frac{2}{5}})y + 5ccx^{-\frac{3}{5}} - 25c^3x^{-\frac{2}{5}}} = \text{Const.}$$

And if in this fraction one puts

$$\text{the numerator} \quad (3 + 15cx^{\frac{1}{5}} + 25ccx^{\frac{2}{5}})y + 5ccx^{-\frac{3}{5}} + 25c^3x^{-\frac{2}{5}} = P \quad \text{and}$$

$$\text{the denominator} \quad (3 - 15cx^{\frac{1}{5}} + 25ccx^{\frac{2}{5}})y + 5ccx^{-\frac{3}{5}} - 25c^3x^{-\frac{2}{5}} = Q,$$

the divisor rendering the propounded equation integrable will be =  $PQ$ .

CASE 6 IN WHICH IT IS  $n = \frac{3}{5}$

§79 Therefore, for this equation

$$dy + yydx - ccx^{-\frac{12}{5}}dx = 0$$

it will be

$$B = \frac{3A}{5c} \quad \text{and} \quad C = \frac{B}{5c} = \frac{3A}{25cc}, \quad D = 0 \quad \text{etc.}$$

and hence the particular integral arises as:

$$y = cx^{-\frac{6}{5}} + \frac{15ccx^{-\frac{2}{5}} + 12cx^{-\frac{1}{5}} + 3}{25ccx^{\frac{3}{5}} + 15cx^{\frac{4}{5}} + 3x}$$

or

$$y = \frac{25c^3x^{-\frac{3}{5}} + 30ccx^{-\frac{2}{5}} + 15cx^{-\frac{1}{5}} + 3}{25ccx^{\frac{3}{5}} + 15cx^{\frac{4}{5}} + 3x} = v,$$

whence the complete integral is obtained:

$$e^{10cx^{-\frac{1}{5}}} \cdot \frac{(3x - 15cx^{\frac{4}{5}} + 25ccx^{\frac{3}{5}})y - 3 + 15cx^{-\frac{1}{5}} - 30ccx^{-\frac{2}{5}} + 25c^3x^{-\frac{3}{5}}}{(3x + 15cx^{\frac{4}{5}} + 25ccx^{\frac{3}{5}})y - 3 - 15cx^{-\frac{1}{5}} - 30ccx^{-\frac{2}{5}} - 25c^3x^{-\frac{3}{5}}} = \text{Const.}$$

And having neglected the exponential factor  $e^{10cx^{-\frac{1}{5}}}$  the product of the numerator and the denominator will yield the divisor, divided by which the propounded equation becomes integrable.

## PROBLEM 12

**§80** While  $i$  denotes any integer number to exhibit the resolution of this equation:

$$dy + yydx - ccx^{\frac{-4i}{2i+1}}dx = 0.$$

## SOLUTION

Therefore, because it is  $n = \frac{i}{2i+1}$ , one will find

$$B = -\frac{(i+1)i}{2(2i+1)c}A$$

$$C = +\frac{(i+2)(i+1)i(i-1)}{2 \cdot 4(2i+1)^2c^2}A$$

$$D = -\frac{(i+3)(i+2)(i+1)i(i-1)(i-2)}{2 \cdot 4 \cdot 6(2i+1)^3c^3}A$$

$$E = +\frac{(i+4)(i+3)(i+2)(i+1)i(i-1)(i-2)(i-3)}{2 \cdot 4 \cdot 6 \cdot 8(2i+1)^4c^4}A$$

etc.,

on the other hand the particular integral will be:

$$y = cx^{\frac{-2i}{2i+1}} + \frac{\frac{i}{2i+1}Ax^{\frac{-i-1}{2i+1}} + \frac{i-1}{2i+1}Bx^{\frac{-i-2}{2i+1}} + \frac{i-2}{2i+1}Cx^{\frac{-i-3}{2i+1}} + \frac{i-3}{2i+1}Dx^{\frac{-i-4}{2i+1}} + \text{etc.}}{Ax^{\frac{i}{2i+1}} + Bx^{\frac{i-1}{2i+1}} + Cx^{\frac{i-2}{2i+1}} + Dx^{\frac{i-3}{2i+1}} + \text{etc.}}$$

to reduce which to the same denominator, let us set:

$$\begin{aligned}\mathfrak{A} &= cA \\ \mathfrak{B} &= \frac{i(i-1)}{2(2i+1)}A \\ \mathfrak{C} &= \frac{(i+1)i(i-1)(i-2)}{2 \cdot 4(2i+1)^2c}A \\ \mathfrak{D} &= \frac{(i+2)(i+1)i(i-1)(i-2)(i-3)}{2 \cdot 4 \cdot 6(2i+1)^3c^2}A \\ &\text{etc.,}\end{aligned}$$

whence it will be:

$$y = \frac{\mathfrak{A}x^{\frac{-i}{2i+1}} + \mathfrak{B}x^{\frac{-i-1}{2i+1}} + \mathfrak{C}x^{\frac{-i-2}{2i+1}} + \mathfrak{D}x^{\frac{-i-3}{2i+1}}}{Ax^{\frac{i}{2i+1}} + Bx^{\frac{i-1}{2i+1}} + Cx^{\frac{i-2}{2i+1}} + Dx^{\frac{i-3}{2i+1}}}.$$

Further, for the sake of brevity let us put:

$$\begin{aligned}Ax^{\frac{i}{2i+1}} + Bx^{\frac{i-1}{2i+1}} + Cx^{\frac{i-2}{2i+1}} + Dx^{\frac{i-3}{2i+1}} + \text{etc.} &= P \\ Ax^{\frac{i}{2i+1}} - Bx^{\frac{i-1}{2i+1}} + Cx^{\frac{i-2}{2i+1}} - Dx^{\frac{i-3}{2i+1}} + \text{etc.} &= Q \\ \mathfrak{A}x^{\frac{-i}{2i+1}} + \mathfrak{B}x^{\frac{-i-1}{2i+1}} + \mathfrak{C}x^{\frac{-i-2}{2i+1}} + \mathfrak{D}x^{\frac{-i-3}{2i+1}} + \text{etc.} &= \mathfrak{P} \\ -\mathfrak{A}x^{\frac{-i}{2i+1}} + \mathfrak{B}x^{\frac{-i-1}{2i+1}} - \mathfrak{C}x^{\frac{-i-2}{2i+1}} + \mathfrak{D}x^{\frac{-i-3}{2i+1}} - \text{etc.} &= \mathfrak{Q}\end{aligned}$$

and the complete integral will be:

$$e^{-2(2i+1)cx^{\frac{i+1}{2i+1}}} \frac{Qy - \mathfrak{Q}}{Py - \mathfrak{P}} = \text{Const.}$$

But then the divisor rendering the propounded equation integrable will be  $(Py - \mathfrak{P})(Qy - \mathfrak{Q})$ .

#### COROLLARY 1

§81 Therefore, if in the equation

$$dy + yydx + ax^{\frac{-4i}{2i+1}}dx = 0$$



the coefficient  $\alpha$  was a negative quantity, such that having put  $\alpha = -cc$   $c$  is a real quantity, the complete integral found here has a real form, and can be easily exhibited in each case, equally as the divisor, which renders the equation integrable.

#### COROLLARY 2

§82 But if  $\alpha$  was a positive quantity, say  $\alpha = aa$  that one has this equation:

$$dy + yydx + aax^{\frac{-4i}{2i+1}}dx = 0,$$

it will be  $c = a\sqrt{-1}$ , and the coefficients  $B, D, F$  etc. and  $\mathfrak{A}, \mathfrak{C}, \mathfrak{E}$  will become negative; hence the particular values  $y = \frac{\mathfrak{P}}{p}$  and  $y = \frac{\mathfrak{Q}}{q}$  will arise as imaginary.

#### COROLLARY 3

§83 Nevertheless in the case, in which it is  $c = a\sqrt{-1}$  and  $cc = -aa$ ,  $P + Q$  and  $\mathfrak{P} + \mathfrak{Q}$  will become real quantities, but  $p - Q$  and  $\mathfrak{P} - \mathfrak{Q}$  imaginary ones. Therefore, if one puts

$$P + Q = 2R, \quad P - Q = 2S\sqrt{-1}, \quad \mathfrak{P} + \mathfrak{Q} = 2\mathfrak{R} \quad \text{and} \quad \mathfrak{P} - \mathfrak{Q} = 2\mathfrak{S}\sqrt{-1},$$

$R, S, \mathfrak{R}$  and  $\mathfrak{S}$  will be real quantities and because of

$$P = R + S\sqrt{-1}, \quad Q = R - S\sqrt{-1}, \quad \mathfrak{P} = \mathfrak{R} + \mathfrak{S}\sqrt{-1}, \quad \mathfrak{Q} = \mathfrak{R} - \mathfrak{S}\sqrt{-1}$$

the divisor rendering the equation integrable will become

$$(RR + SS)yy - 2(R\mathfrak{R} + S\mathfrak{S})y + \mathfrak{R}\mathfrak{R} + \mathfrak{S}\mathfrak{S}$$

and hence real.

#### COROLLARY 4

§84 But in the same case  $c = a\sqrt{-1}$  because of

$$e^{-p\sqrt{-1}} = \cos p - \sqrt{-1} \sin p,$$

it will be

$$e^{-2(2i+1)\alpha x^{\frac{1}{2i+1}}\sqrt{-1}} = \cos 2(2i+1)\alpha x^{\frac{1}{2i+1}} - \sqrt{-1} \sin 2(2i+1)\alpha x^{\frac{1}{2i+1}}$$

whence having put for the sake of brevity

$$2(2i+1)\alpha x^{\frac{1}{2i+1}} = p,$$

the complete integral will be:

$$(\cos p - \sqrt{-1} \sin p) \cdot \frac{(R + S\sqrt{-1})y - \mathfrak{R} + \mathfrak{S}\sqrt{-1}}{(R - S\sqrt{-1})y - \mathfrak{R} - \mathfrak{S}\sqrt{-1}} = \text{Const.},$$

which form is imaginary.

#### COROLLARY 5

§85 But attribute such a form to the constant:  $\alpha - \beta\sqrt{-1}$ , and having expanded the integral equation, it will be:

$$\begin{aligned} & (Ry - \mathfrak{R}) \cos p - (Ry - \mathfrak{R}) \sin p \sqrt{-1} - (Sy - \mathfrak{S}) \cos p \sqrt{-1} - (Sy - \mathfrak{S}) \sin p \\ & = (Ry - \mathfrak{R})\alpha - (Ry - \mathfrak{R})\beta\sqrt{-1} + (Sy - \mathfrak{S})\alpha\sqrt{-1} + (Sy - \mathfrak{S})\beta. \end{aligned}$$

Now equate the real and imaginary parts separately:

$$\begin{aligned} (Ry - \mathfrak{R}) \cos p - (Sy - \mathfrak{S}) \sin p &= \alpha(Ry - \mathfrak{R}) + \beta(Sy - \mathfrak{S}) \\ (Ry - \mathfrak{R}) \sin p + (Sy - \mathfrak{S}) \cos p &= \beta(Ry - \mathfrak{R}) - \alpha(Sy - \mathfrak{S}), \end{aligned}$$

which two equations agree, if it only is

$$\alpha\alpha + \beta\beta = 1.$$

Therefore, let  $\alpha = \cos \zeta$  and  $\beta = \sin \zeta$  and from each of them it will arise

$$\frac{Ry - \mathfrak{R}}{Sy - \mathfrak{S}} = \frac{\sin p + \sin \zeta}{\cos p + \cos \zeta} = \cot \frac{\zeta - p}{2}.$$

### COROLLARY 6

§86 Therefore, having taking any angle for  $\zeta$ , if it is  $c = a\sqrt{-1}$ , the complete integral of the propounded equation will be

$$\frac{Ry - \mathfrak{R}}{Sy - \mathfrak{S}} = \cot \frac{\zeta - p}{2}$$

or

$$y = \frac{\mathfrak{R} \sin \frac{\zeta - p}{2} - \mathfrak{S} \cos \frac{\zeta - p}{2}}{R \sin \frac{\zeta - p}{2} - S \cos \frac{\zeta - p}{2}}$$

while  $p = 2(2i + 1)\alpha x^{\frac{1}{2i+1}}$ .

### PROBLEM 13

§87 While  $i$  denotes any arbitrary integer number to exhibit the resolution of this equation:

$$dy + yydx - ccx^{\frac{-4i}{2i-1}}dx = 0.$$

### SOLUTION

Since it is  $n = \frac{i}{2i-1}$ , this resolution can be derived from the solution of the preceding problem by putting  $-i$  instead of  $i$ . Hence, attribute the following values to the letters  $B, C, D$  etc.:

$$\begin{aligned} B &= + \frac{i(i-1)}{2(2i-1)c} A \\ C &= + \frac{(i+1)i(i-1)(i-2)}{2 \cdot 4(2i-1)^2 c^2} A \\ D &= + \frac{(i+2)(i+1)i(i-1)(i-2)(i-3)}{2 \cdot 4 \cdot 6(2i-1)^3 c^3} A \\ &\text{etc.} \end{aligned}$$

But the determination of the other letters  $\mathfrak{A}, \mathfrak{B}, \mathfrak{C}, \mathfrak{D}$  will behave as this:

$$\begin{aligned}\mathfrak{A} &= cA \\ \mathfrak{B} &= + \frac{(i+1)i}{2(2i-1)}A \\ \mathfrak{C} &= + \frac{(i+2)(i+1)i(i-1)}{2 \cdot 4(2i-1)^2c}A \\ \mathfrak{D} &= + \frac{(i+3)(i+2)(i+1)i(i-1)(i-2)}{2 \cdot 4 \cdot 6(2i-1)^3c^2}A \\ &\text{etc.}\end{aligned}$$

Having constituted these values for the sake of brevity put:

$$\begin{aligned}Ax^{\frac{i}{2i-1}} + Bx^{\frac{i+1}{2i-1}} + Cx^{\frac{i+2}{2i-1}} + Dx^{\frac{i+3}{2i-1}} + \text{etc.} &= P \\ Ax^{\frac{i}{2i-1}} - Bx^{\frac{i+1}{2i-1}} + Cx^{\frac{i+2}{2i-1}} - Dx^{\frac{i+3}{2i-1}} + \text{etc.} &= Q \\ \mathfrak{A}x^{\frac{-i}{2i-1}} + \mathfrak{B}x^{\frac{-i+1}{2i-1}} + \mathfrak{C}x^{\frac{-i+2}{2i-1}} + \mathfrak{D}x^{\frac{-i+3}{2i-1}} + \text{etc.} &= \mathfrak{P} \\ -\mathfrak{A}x^{\frac{-i}{2i-1}} + \mathfrak{B}x^{\frac{-i+1}{2i-1}} - \mathfrak{C}x^{\frac{-i+2}{2i-1}} + \mathfrak{D}x^{\frac{-i+3}{2i-1}} - \text{etc.} &= \mathfrak{Q}\end{aligned}$$

and hence one immediately has two particular integrations:

$$\text{I. } y = \frac{\mathfrak{P}}{P} \quad \text{and} \quad \text{II. } y = \frac{\mathfrak{Q}}{Q}.$$

But then the complete integral equation will be:

$$e^{2(2i-1)cx^{\frac{-1}{2i-1}}} \frac{Qy - \mathfrak{Q}}{Py - \mathfrak{P}} = \text{Const.}$$

and the divisor rendering the propounded equation integrable will be =  $(Py - \mathfrak{P})(Qy - \mathfrak{Q})$ .

#### COROLLARY 1

§88 But if the propounded equation was of this kind:

$$dy + yydx + aax^{\frac{-4i}{2i-1}}dx = 0,$$

that it is  $cc = -aa$  and  $c = a\sqrt{-1}$ , the exhibited particular integrations will become imaginary because of the imaginary  $B, D, F$  and  $\mathfrak{A}, \mathfrak{C}, \mathfrak{E}$  etc., whereas the values of the remaining letters are real.

## COROLLARY 2

§89 But if one puts

$$P + Q = 2R, \quad P - Q = 2S\sqrt{-1}, \quad \mathfrak{P} + \mathfrak{Q} = 2\mathfrak{R} \quad \text{and} \quad \mathfrak{P} - \mathfrak{Q} = 2\mathfrak{S}\sqrt{-1},$$

the quantities  $R, S, \mathfrak{R}$  and  $\mathfrak{S}$  will nevertheless, as before, be real, and the divisor rendering the equation integrable will be:

$$(RR + SS)yy - 2(R\mathfrak{R} + S\mathfrak{S})y + \mathfrak{R}\mathfrak{R} + \mathfrak{S}\mathfrak{S}.$$

## COROLLARY 3

§90 But then, of one for the sake of brevity puts

$$2(2i - 1)\alpha x^{\frac{-i}{2i-1}} = p,$$

the complete integral will be:

$$\frac{Ry - \mathfrak{R}}{Sy - \mathfrak{S}} = \cot \frac{\zeta + p}{2},$$

whence one finds:

$$y = \frac{\mathfrak{R} \sin \frac{\zeta + p}{2} - \mathfrak{S} \cos \frac{\zeta + p}{2}}{R \sin \frac{\zeta + p}{2} - S \cos \frac{\zeta + p}{2}}$$

where the angle  $\zeta$  takes the part of the arbitrary constant.

## SCHOLIUM

§91 The solutions of these last two problems were not expanded so by accurate analysis as derived by induction from the particular cases explained above, since the progression from these cases to the following was sufficiently manifest. But the foundation of these solution mainly lies in this, that a particular solution, whence all are deduced is actually a double one, since the quantity  $c$ , only whose square occurs in the differential equation, can be taken negatively and positively. But as often as two particular solutions of equations of this kind are known, from them the general solution and hence the multipliers rendering it integrable can be found a lot easier, which will be worth one's while to have explained it more clearly.

PROBLEM 14

§92 Having found to particular solutions of an equation of this kind:

$$dy + Pydx + Qyydx + Rdx = 0$$

to find its general solution and multiplicator which renders it integrable.

SOLUTION

Let  $M$  and  $N$  be functions of  $x$  of such a kind, which substituted instead of  $y$  satisfy the propounded equation such that it is:

$$dM + PMdx + QM^2dx + Rdx = 0$$

and

$$dN + PNdx + QN^2dx + Rdx = 0.$$

Put

$$\frac{y - M}{y - N} = z \quad \text{and} \quad y = \frac{M - Nz}{1 - z},$$

it will be

$$dy = \frac{dM - zdM + Mdz - Ndz - zdN + zzdN}{(1 - z)^2},$$

having substituted these values in the propounded equation and having multiplied the whole equation by  $(1 - z)^2$  it will arise:

$$(1 - z)dM - z(1 - z)dN + (M - N)dz + P(1 - z)Mdx - P(1 - z)Nzdx \\ + QMMdx - 2QMNzdx + QNNzzdx + R(1 - z)^2dx = 0.$$

Now substitute the values to arise from the two superior differentials for  $dM$  and  $dN$ :

$$\begin{aligned}
& -P(1-z)Mdx - Q(1-z)M^2dx - R(1-z)dx \\
& + Pz(1-z)Ndx + Qz(1-z)N^2dx + Rz(1-z)dx + (M-N)dx = 0 \\
& + P(1-z)Mdx + QM^2dx + R(1-z)^2dx \\
& - Pz(1-z)Ndx - 2QMNzdx \\
& + QN^2zdx,
\end{aligned}$$

having ordered which equation it will arise:

$$QzM^2dx + QzN^2dx - 2QMNzdx + (M-N)dz = 0$$

or

$$Q(M-N)dx + \frac{dz}{z} = 0,$$

such that it is:

$$z = Ce^{-\int Q(M-N)dx},$$

whence the general integrated equation will be:

$$e^{\int Q(M-N)dx} \frac{y-M}{y-N} = \text{Const.}$$

But for finding the multiplier note that the propounded equation having done the substitution at first was multiplied by  $(1-z)^2$ , but then divided by  $z(M-N)$  became integrable. Therefore, it multiplied  $\frac{(1-z)^2}{(M-N)z}$  immediately will become integrable: from this the factor will be  $\frac{(1-z)^2}{(M-N)z}$ , which because of  $z = \frac{y-M}{y-N}$  will obtain this form:

$$\frac{M-N}{(y-M)(y-N)}.$$

#### PROBLEM 15

§93 Having propounded the equation

$$ydy + Pydx + Qdx = 0,$$

to find conditions of the functions  $P$  and  $Q$  that a multiplier of this kind  $(y+M)^n$  renders it integrable.

### SOLUTION

Therefore, from the nature of differentials it must be:

$$\frac{1}{dx}d.y(y + M) = \frac{1}{dy}d.(Py + Q)(y + M)^n,$$

because hence  $M$  is a function of  $x$  only, it will be

$$ny(y + M)^{n-1}\frac{dM}{dx} = P(y + M)^n + n(Py + Q)(y + M)^{n-1},$$

which divided by  $(y + M)^{n-1}$  goes over into this one:

$$\frac{nydM}{dx} = (n + 1)Py + PM + nQ,$$

whence it is necessary that it is:

$$P = \frac{ndM}{(n + 1)dx} \quad \text{and} \quad Q = \frac{-PM}{n} = -\frac{MdM}{(n + 1)dx}.$$

Therefore, having substituted these values the equation

$$ydy + \frac{nydM}{n + 1} - \frac{MdM}{n + 1} = 0$$

becomes integrable, if it is multiplied by  $(y + M)^n$ .

### COROLLARY 1

§94 Since this equation is homogeneous, it is also integrable, if it is divided by

$$(n + 1)yy + nyM - MM = (y + M)((n + 1)y - M).$$

And therefore hence no new equations treatable by this method are obtained.

### COROLLARY 2

§95 But since we have these two multipliers

$$(y + M)^n \quad \text{and} \quad \frac{1}{(y + M)((n + 1)y - M)},$$



if the one is divided by the other, the quotient equated to an arbitrary constant will give the complete integral. Hence the equation

$$ydy + \frac{nydM}{n+1} - \frac{MdM}{n+1} = 0$$

integrated generally yields:

$$(y + M)^{n+1}((n + 1)y - M) = \text{Const.}$$

#### PROBLEM 16

§96 Having propounded the equation

$$ydy + Pydx + Qdx = 0,$$

to find the conditions of the functions  $P$  and  $Q$  that a multiplier of this kind

$$(yy + My + N)^n$$

renders it integrable.

#### SOLUTION

From the nature of differentials it is necessary that it is:

$$\frac{1}{dx}d.y(yy + My + N)^n = \frac{1}{dy}d.(Py + Q)(yy + My + N)^n.$$

Therefore, since  $M$ ,  $N$ ,  $P$  and  $Q$  by assumption are functions of  $x$ , it will be having done the expansion:

$$\begin{aligned} & ny(yy + My + N)^{n-1} \left( y + \frac{dM}{dx} + \frac{dN}{dx} \right) \\ & = P(yy + My + N)^n + n(Py + Q)(1y + M)(yy + My + N)^{n-1} \end{aligned}$$

and after division by  $(yy + My + N)^{n-1}$ :

$$\begin{aligned} nyy\frac{dM}{dx} + \frac{nydN}{dx} &= (2n + 1)Py + (n + 1)PM + PN \\ & \quad + 2nQy \quad + nQM. \end{aligned}$$

Hence it must be:

- I.  $ndM = (2n + 1)Pdx$
- II.  $(n + 1)PMdx + 2nQdx$
- III.  $0 = PN + nQM.$

The first gives

$$P = \frac{ndM}{(2n + 1)dx}$$

and the last

$$Q = \frac{-PN}{nM} \quad \text{or} \quad Q = \frac{-NdM}{(2n + 1)Mdx},$$

which values substituted in the middle one yield:

$$ndN = \frac{n(N + 1)MdM}{2n + 1} - \frac{2nNdM}{(2n + 1)M}$$

or

$$(2n + 1)MdN + 2NdM = (n + 1)MMdM,$$

which multiplied by  $M^{\frac{-2n+1}{2n+1}}$  and integrated yields:

$$(2n + 1)M^{\frac{2}{2n+1}}N = \text{Const.} + (n + 1) \int M^{\frac{2n+3}{2n+1}} dM$$

or

$$(2n + 1)M^{\frac{2}{2n+1}}N = \text{Const.} + \frac{2n + 1}{4} M^{\frac{4n+4}{2n+1}},$$

whence it is

$$N = \alpha M^{\frac{-2}{2n+1}} + \frac{1}{4} M^2.$$

Therefore, because it is

$$Pdx = \frac{ndM}{2n + 1} \quad \text{and} \quad Qdx = -\frac{\alpha M^{\frac{-2n-3}{2n+1}} dM}{2n + 1} - \frac{MdM}{4(2n + 1)},$$

this differential equation:

$$ydy + \frac{nydM}{2n+1} - \frac{MdM}{4(2n+1)} - \frac{\alpha}{2n+1} M^{\frac{-2n-3}{2n+1}} dM = 0$$

is rendered integrable, if it is multiplied by

$$\left( yy + My + \frac{1}{4}M^2 + \alpha M^{\frac{-2}{2n+1}} \right)^n.$$

#### COROLLARY 1

§97 If it was

$$\frac{-2n-3}{2n+1} = 1 \quad \text{or} \quad n = -1,$$

the differential equation is homogeneous, and if

$$\frac{-2n-3}{2n+1} = 0 \quad \text{or} \quad n = -\frac{3}{2},$$

it is of degree one. But in each of both cases there is no difficulty, since the equation can easily be treated.

#### COROLLARY 2

§98 Therefore, the cases, in which the exponent  $\frac{-2n-3}{2n+1}$  is neither 0 nor 1, will be more strange. Therefore, let

$$\frac{-2n-3}{2n+1} = m, \quad \text{whence it is} \quad 2n = \frac{-m-3}{m+1},$$

and the differential equation

$$ydy + \frac{1}{4}(m+3)ydM + \frac{1}{8}(m+1)MdM + \frac{1}{2}\alpha(m+1)M^m dM = 0$$

will be rendered integrable by the multiplier

$$(yy + My + \frac{1}{4}MM + \alpha M^{m+1})^{\frac{-m-3}{2(m+1)}}.$$

### COROLLARY 3

§99 If now for  $M$  any functions of  $x$  are substituted, one will be able to form such complicated equations, how which have to treated by other methods is hardly clear, although by this method their resolution is obvious.

### SCHOLIUM

§100 I anyone wants to follow this path further, there is no doubt that this method will soon obtain a lot greater increments, by which the whole field of Analysis is significantly promoted. The specimens expanded here are also of such a nature that they seem to pave the way to more profound investigations, especially if additionally other classes of differential equations are treated in similar manner. But these things, I presented up to now, seem to suffice to encourage the Geometers to develop this method further, which goal I had mainly set myself.