

# **Ultraviolet Spectroscopy of the Actinium-229 beta decay: On the way to the first identification of the $^{229}\text{Th}$ low-energy isomer?**

**S. Kraemer<sup>1</sup> for the ISOLDE-IS658 collaboration**

**<sup>1</sup>Instituut voor Kern- en Stralingsfysica, KU Leuven**

A unique feature of thorium-229 is its isomer with an exceptionally low excitation energy, proposed as a candidate for future optical clocks [1]. The small decay width is expected to outperform the accuracy of current state-of-the-art atomic clocks by an order of magnitude [2]. The current best values of the excitation energy are 8.28(17)eV and 8.10(17)eV [3,4]. These were determined using two different measurement techniques whereby the isomer is populated in the alpha decay of uranium-233. The development of an optical clock requires however knowledge of the excitation energy by at least an order of magnitude more precise. Spectroscopic experiments searching for a direct signature of the radiative decay have to-date been unsuccessful, partially due to the background induced in the preceding alpha decay.

An alternative approach using the beta decay of actinium-229 is studied as a novel method to populate the isomer with high efficiency and in low background conditions [5]. Produced online at the ISOLDE facility, actinium is laser-ionized and implanted into a large-bandgap crystal in specific lattice positions, suppressing the electron conversion decay channel of the isomer. A favourable feeding pattern is significantly increasing the population of the isomer compared to uranium-233 and the lower energy deposit of the beta compared to the alpha decay results in a significantly reduced luminescence background.

In this seminar, a brief overview over the topic is given, a dedicated setup developed at KU Leuven for the implantation of an actinium-229 beam into large-bandgap crystals and the vacuum-ultraviolet spectroscopic study of the emitted photons is presented and the preliminary results from a recent experimental campaign at ISOLDE are presented.

[1] E. Peik et al., *Europhys. Lett.* 61, 2 (2003)

[2] C. Campbell et al., *PRL* 108, 120802 (2012)

[3] B. Seiferle et al., *Nature* 573, 243-246 (2019)

[4] T. Sikorsky et al., *PRL* 125, 142503 (2020)

[5] M. Verlinde et al., *Physical Review C*, 100 (2), 024315-024315