

# ON THE USE OF DIFFERENTIAL CALCULUS IN THE RESOLUTION OF EQUATIONS \*

Leonhard Euler

§227 That the constitution of an equation can be reduced to the nature of functions was shown sufficiently enough above already. For, let  $y$  denote any function of  $x$ ; if one puts  $y = 0$ , in this form completely all finite equations, may they be algebraic or transcendental, are contained. But the equation  $y = 0$  is said to be solved, if that value  $x$  is attributed to it which substituted in the function  $y$  actually renders it equal to nothing. But in most cases many of such values are given for  $x$ , which are called the roots of the equation  $y = 0$ . Therefore, if we put that the numbers  $f, g, h, i$  etc. are roots of the equation  $y = 0$ , the function  $y$  will be of such a nature that, if in it instead of  $x$  either  $f$  or  $g$  or  $h$  or etc., is substituted, it indeed is  $y = 0$ .

§227 Therefore, since the function  $f$  vanishes, if in it instead of  $x$  one puts  $f$  or  $x + (f - x)$ , where  $f$  is a root of the equation  $y = 0$ , it will be by means of that what we demonstrated about functions above [§48],

$$0 = y + \frac{(f - x)dy}{dx} + \frac{(f - x)^2 ddy}{2dx^2} + \frac{(f - x)^3 d^3y}{6dx^3} + \text{etc.},$$

from which equation the value of the root  $f$  is determined in such a way that, whatever was put for  $x$  and hence the value of the quantities  $y, \frac{dy}{dx}, \frac{ddy}{2dx^2}$  etc.

---

\*Original title: "De Usu Calculi Differentialis in Aequationibus resolvendis", first published as part of the book „*Institutiones calculi differentialis cum eius usu in analysi finitorum ac doctrina serierum*, 1755“, reprinted in in „*Opera Omnia: Series 1, Volume 10*, pp. 422 - 445“, Eneström-Number E212, translated by: Alexander Aycocock for the „Euler-Kreis Mainz“

were substituted, always the equation expressing the true value of  $f$  results. That this is seen more clearly, let us put that it is

$$y = x^3 - 2x^2 + 3x - 4;$$

it will be

$$\frac{dy}{dx} = 3x^2 - 4x + 3, \quad \frac{d^2y}{dx^2} = 3x - 2, \quad \text{and} \quad \frac{d^3y}{dx^3} = 1.$$

Having substituted these values it arises

$$0 = x^3 - 2x^2 + 3x - 4 + (f - x)(3x^2 - 4x + 3) + (f - x)^2(3x - 2) + (f - x)^3$$

or having actually done the multiplications

$$f^3 - 2ff + 3f - 4 = 0;$$

of course the same equation as the propounded one arises which therefore contains the same roots.

**§229** But although this way one does not get to a new equation, from which the value of the root  $f$  can be determined in an easier way, nevertheless from this extraordinary auxiliary theorems for the invention of roots can be deduced. For, if for  $x$  a value already very close to a certain root was assumed such that  $f - x$  is a very small quantity, then the terms of the equation

$$0 = y + \frac{(f - x)dy}{dx} + \frac{(f - x)^2 d^2y}{2dx^2} + \frac{(f - x)^3 d^3y}{6dx^3} + \text{etc.}$$

will converge very rapidly and therefore one will not deviate much from the truth, if except the first two initial terms all remaining ones are rejected. Therefore, it will be, if for  $x$  a value already close to a certain root of the equation  $y = 0$  was assumed, approximately

$$0 = y + \frac{(f - x)dy}{dx} \quad \text{or} \quad f = x - \frac{ydx}{dy},$$

from which formula an, even though not true, but nevertheless very good approximate value of the root  $f$  will be found, which thereafter again substituted for  $x$  will yield a even better value for  $f$  and so on will continuously get closer to the true value of the root  $f$ .

§230 Therefore, at first the root of all powers of any number can be taken. For, let the number  $a^n + b$  be propounded, of which the root of the power  $n$  shall to be extracted. Put  $x^n = a^n + b$  or  $x^n - a^n - b = 0$  that it is  $y = x^n - a^n - b$ ; it will be

$$\frac{dy}{dx} = nx^{n-1}, \quad \frac{ddy}{2dx^2} = \frac{n(n-1)}{1 \cdot 2}x^{n-2}, \quad \frac{d^3y}{6dx^3} = \frac{n(n-1)(n-2)}{1 \cdot 2 \cdot 3}x^{n-3} \quad \text{etc.}$$

Hence, if the root in question is put =  $f$  that it is  $f = \sqrt[n]{a^n + b}$ , it will be

$$0 = x^n - a^n - b + n(f-x)x^{n-1} + \frac{n(n-1)}{1 \cdot 2}(f-x)^2x^{n-2} + \text{etc.}$$

Therefore, if for  $x$  one sets a number already coming close to the value of the root  $f$  in question, which will happen by putting  $x = a$ , if  $b$  was so a small number that  $a^n + b < (a+1)^n$ , it will approximately be  $b = na^{n-1}(f-a)$  and hence

$$f = a + \frac{b}{na^{n-1}},$$

whence the value of the root will be known a lot closer. But if we want to take also the third term that it is

$$b = na^{n-1}(f-a) + \frac{n(n-1)}{1 \cdot 2}a^{n-2}(f-a)^2,$$

it will be

$$(f-a)^2 = -\frac{2a}{n-1}(f-a) + \frac{2b}{n(n-1)a^{n-2}}$$

and hence

$$f = a - \frac{a}{n-1} \pm \sqrt{\frac{aa}{(n-1)^2} + \frac{2b}{n(n-1)a^{n-2}}}$$

or

$$f = \frac{(n-2)a + \sqrt{aa + 2(n-1)b : na^{n-2}}}{n-1}.$$

Hence, by means of the extraction of the square root the value of the root  $f$  will be found even closer.

EXAMPLE

Let us find the square root of any number  $c$  or let  $xx - c = y$ .

Therefore, put the number very close to the root =  $a$  and  $b = c - aa$ ; because of  $aa + b = c$  and since it is  $n = 2$ , the first formula will become  $f = a + \frac{c-aa}{2a} = \frac{c+aa}{2a}$ ; the other gives  $f = \sqrt{c}$  which is the root in question itself. Therefore, because the root approximately is =  $\frac{c+aa}{2a}$ , write the value itself for  $a$  and a closer root will be  $f = \frac{cc+6aac+a^4}{4a(c+aa)}$ . For the sake of an example let  $c = 5$ , from the first formula it will be  $f = \frac{5}{2a} + \frac{a}{2}$ . Therefore, put  $a = 2$ , it will be  $f = 2.25$ ; now put  $a = 2.25$ , it will be  $f = 2.236111$ ; further, set  $a = 2.236111$ , it will be  $f = 2.2360679$  which value already hardly deviated from the truth.

§231 But in similar matter the root of any equation can be found approximately by means of the equation  $f = x - \frac{ydx}{dy}$ , after having assumed a value differing hardly from a certain root of the equation for  $x$ , of course. To find a value of this kind for  $x$ , successively put various values for  $x$  and from them chose the one which minimizes the function  $y$ , which means indicates the value closest to zero. So, if it is

$$y = x^3 - 2xx + 3x - 4$$

having put	$x = 0$	it is	$y = -4$
	$x = 1$		$y = -2$
	$x = 2$		$y = +2,$

whence we see that the root is contained within the values 1 and 2 of  $x$ . Therefore, because it is  $\frac{dy}{dx} = 3xx - 4x + 3$ , for finding the root  $f$  of the equation  $x^3 - 2xx + 3x - 4 = 0$  one will have this equation

$$f = x - \frac{ydx}{dy} = x - \frac{x^3 - 2xx + 3x - 4}{3xx - 4x + 3}.$$

Therefore, let  $x = 1$ ; it will be  $f = 1 + \frac{2}{2} = 2$ . Now, put  $x = 2$ ; it will be  $f = 2 - \frac{2}{7} = \frac{12}{7}$ . Therefore, let  $x = \frac{12}{7}$ ; it will be  $f = \frac{12}{7} - \frac{104}{1701} = \frac{2812}{1701} = 1.653$ . If we want to proceed further, we will use logarithms more conveniently. Therefore, put  $x = 1.653$  and it will be

$\log x^1$	$=$	0.2182729	$x^1$	$=$	1.653000
$\log x^2$	$=$	0.4365458	$x^2$	$=$	2.732409
$\log x^3$	$=$	0.6548187	$x^3$	$=$	4.516673
$x^3$	$=$	4.516673			
$3x$	$=$	<u>4.959000</u>			

And hence

$x^3 + 3x$	$=$	9.475673	$3xx + 3$	$=$	11.197227
$2xx + 4$	$=$	<u>9.464818</u>	$4x$	$=$	<u>6.612000</u>
num.	$=$	0.010855	den.	$=$	4.585227
$\log$ num.	$=$	8.0356298			
$\log$ den.	$=$	0.6613608	$x$	$=$	1.6553000
$\log$ fract.	$=$	7.3742690	fraction	$=$	<u>0.002367</u>
			$f$	$=$	<u>1.650633,</u>

which value already comes very close to the true one.

§232 But we will be able to deduce faster approximations from the general expression. For, because having put any function  $y = 0$ , if the root of this equation was  $x = f$ , we will find that it is

$$0 = y + \frac{(f-x)dy}{dx} + \frac{(f-x)^2 ddy}{2dx^2} + \frac{(f-x)^3 d^3y}{6dx^3} + \text{etc.},$$

let  $f - x = z$ , such that the root is  $f = x + z$ , and put

$$\frac{dy}{dx} = p, \quad \frac{dp}{dx} = q, \quad \frac{dq}{dx} = r, \quad \frac{dr}{dx} = s \quad \text{etc.};$$

it will be

$$0 = y + zp + \frac{z^2q}{2} + \frac{z^3r}{6} + \frac{z^4s}{24} + \frac{z^5t}{120} + \text{etc.};$$

in this equation having taken any value for  $x$ , from which at the same time  $y, p, q, r, s$  etc. are determined, the quantity  $z$  must be found, having found which one will have the root  $f = x + z$  of the propounded equation  $y = 0$ . Therefore, will have to focus on the task to find the value of the unknown  $z$  from this equation in the most convenient way possible.

§233 Assume this convergent series for  $z$

$$z = A + B + C + D + E + \text{etc.}$$

and having done the substitution it will be

$$\begin{aligned}
 y &= y \\
 pz &= Ap + Bp + Cp + Dp + Ep + \text{etc.} \\
 \frac{1}{2}qz^2 &= \frac{1}{2}A^2q + ABq + ACq + ADq + \text{etc.} \\
 &\quad + \frac{1}{2}BBq + BCq + \text{etc.} \\
 \frac{1}{6}rz^3 &= \frac{1}{6}A^3r + \frac{1}{2}A^2Br + \frac{1}{2}A^2Cr + \text{etc.} \\
 &\quad + \frac{1}{2}AB^2r + \text{etc.} \\
 \frac{1}{24}sz^4 &= \frac{1}{24}A^4s + \frac{1}{6}A^3Bs + \text{etc.} \\
 \frac{1}{120}tz^5 &= \frac{1}{120}A^5t + \text{etc.}
 \end{aligned}$$

Hence, one will obtain the following equation

$$\begin{aligned}
A &= -\frac{y}{p} \\
B &= -\frac{yyq}{2p^3} \\
C &= -\frac{y^3qq}{2p^5} + \frac{y^3r}{6p^4} \\
D &= -\frac{5y^4q^3}{8p^7} + \frac{5y^4qr}{12p^6} - \frac{y^4s}{24p^5} \\
&\text{etc.}
\end{aligned}$$

and hence it will be

$$z = -\frac{y}{p} - \frac{y^2q}{2p^3} - \frac{y^3qq}{2p^5} + \frac{y^3r}{6p^4} - \frac{5y^4q^3}{8p^7} + \frac{5y^4qr}{12p^6} - \frac{y^4s}{24p^5} - \text{etc.}$$

#### EXAMPLE

Let this equation be propounded  $x^5 + 2x - 2 = 0$ .

Therefore, it will be

$$\begin{aligned}
y = x^5 + 2x - 2, \quad \frac{dy}{dx} = p = 5x^4 + 2, \quad \frac{dp}{dx} = q = 20x^3, \\
\frac{dq}{dx} = r = 60x^2, \quad \frac{dr}{dx} = s = 120x \quad \text{etc.}
\end{aligned}$$

But now put  $x = 1$ , since this value hardly deviates from the root, it will be

$$y = 1, \quad p = 7, \quad q = 20, \quad r = 60, \quad s = 120,$$

whence it will be

$$z = -\frac{1}{7} - \frac{10}{7^3} - \frac{200}{7^5} + \frac{10}{7^4} - \frac{5 \cdot 1000}{7^7} + \frac{500}{7^6} - \frac{5}{7^5} + \text{etc.}$$

or

$$z = -\frac{1}{7} - \frac{10}{7^3} - \frac{130}{7^5} - \frac{1745}{7^7} - \text{etc.},$$

and therefore it will be  $z = 0.18$  and the root  $f = 0.82$ ; if this value is again substituted instead of  $x$ , a root very close to the true one will arise.

§234 Therefore, we found an infinite series, which expresses the root of any equation; but it has this inconvenience that then the law of progression is not plain, and it hence is too complex and not sufficiently useful. Therefore, we undertake the same task in another way and investigate a more regular series expressing any root of the propounded equation.

Let as before the equation  $y = 0$  be propounded while  $y$  is any function of  $x$  and the question reduces to that the value of  $x$  is defined which substituted for  $x$  renders the function  $y$  equal to zero. But because  $y$  is a function of  $x$ , vice versa  $x$  can be considered as function of  $y$  and considering it like this the value of the function  $x$  is to be found which it takes, if the quantity  $y$  vanishes. Therefore, if it is propounded to give the value of  $x$  which will be the root of the equation  $y = 0$ , since  $x$  goes over into  $f$ , if one sets  $y = 0$ , it will be by means of that what was demonstrated above [§ 67],

$$f = x - \frac{ydx}{dy} + \frac{y^2ddx}{2dy^2} - \frac{y^3d^3}{6dy^3} + \frac{y^4d^4x}{24dy^4} - \text{etc.},$$

in which equation the differential  $dy$  is set constant. Therefore, if one puts

$$\frac{dx}{dy} = p, \quad \frac{dp}{dy} = q, \quad \frac{dq}{dy} = r, \quad \frac{dr}{dy} = s \quad \text{etc.},$$

having introduced these values, that the consideration of a constant differential is avoided, it will be

$$f = x - py + \frac{1}{2}qy^2 - \frac{1}{6}ry^3 + \frac{1}{24}sy^4 - \frac{1}{120}ty^5 + \text{etc.}$$

§235 Therefore, having attributed any value to  $x$  at the same time the values of  $y$  and the quantities  $p, q, r, s$  etc. will be determined and having found these values one will have an infinite series expressing the value of the root  $f$ . But if the equation  $y = 0$  has several roots, then they arise, if different values are assumed for  $x$ ; for, because  $y$  can take the same value, even though different values are attributed to  $x$ , it is not unexpected that the same series can often yield several values. To avoid this ambiguity in these cases and to render the series convergent, for  $x$  a value already close to the value of its root, which is in question, must be assumed. For, this way the value of  $y$  will become very small and the terms of the series will decrease immensely, such that by taking only a few terms one will already find a sufficiently correct value for  $f$ . If then this value is substituted for  $x$ , the quantity  $y$  will become a lot smaller



and the series will converge a lot more and this way the root  $f$  immediately becomes known so accurately that the error will be very small. And hence the advantages of this expression over the one we found before is manifestly seen.

**§236** Let us put that the root of the power  $n$  of any number  $N$  is to be extracted. Therefore, having taken an approximate power of the exponent  $n$  the propounded number will easily be resolved into this for  $N = a^n + b$ . Therefore, it will be

$$x^n = a^n + b \quad \text{and} \quad y = x^n - a^n - b,$$

whence it is

$$\begin{aligned} dy &= nx^{n-1}dx & \text{and} \quad \frac{dx}{dy} = p &= \frac{1}{nx^{n-1}} \\ dp &= -\frac{(n-1)dx}{nx^n} & \text{and} \quad \frac{dp}{dy} = q &= -\frac{n-1}{nmx^{2n-1}} \\ dq &= \frac{(n-1)(2n-1)dx}{nmx^{2n}} & \text{and} \quad \frac{dq}{dy} = r &= \frac{(n-1)(2n-1)}{n^3x^{3n-1}} \\ dr &= -\frac{(n-1)(2n-1)(3n-1)dx}{n^3x^{3n}} & \text{and} \quad \frac{dr}{dy} = s &= -\frac{(n-1)(2n-1)(3n-1)}{n^4x^{4n-1}} \\ & & & \text{etc.} \end{aligned}$$

Now, put  $x = a$  and it will be  $y = -b$  and the root in question  $f = \sqrt[n]{a^n + b}$  will be expressed this way

$$f = a + \frac{b}{na^{n-1}} - \frac{(n-1)bb}{n \cdot 2na^{2n-1}} + \frac{(n-1)(2n-1)b^3}{n \cdot 2n \cdot 3na^{3n-1}} - \frac{(n-1)(2n-1)(3n-1)b^4}{n^4 \cdot 2n \cdot 3n \cdot 4na^{4n-1}} + \text{etc.}$$

and so the same series arises which is usually found by expansion of the binomial  $(a^n + b)^{\frac{1}{n}}$ .

**§237** Therefore, after in the actual extraction the approximate root  $a$  was found and at the same time the residue  $b$  was found then to the root additionally the value  $\frac{b}{an^{n-1}}$  is to be added, that a root closer to the true one is obtained. But it will be

$$a^{n-1} = \frac{N - b}{a}$$

because of  $N = a^n + b$ . But this way a root larger than the correct one will be found, since the third term must be subtracted. Therefore, to find a root a lot closer to the true one by means of division of the residue  $b$ , a suitable divisor must be investigated, which shall be assumed to be

$$na^{n-1} + \alpha b + \beta b^2 + \gamma b^3 + \text{etc.}$$

Therefore, because it must be

$$\begin{aligned} & \frac{b}{na^{n-1} + \alpha b + \beta b^2 + \gamma b^3 + \text{etc.}} \\ = & \frac{b}{na^{n-1}} - \frac{(n-1)bb}{2n^2a^{2n-1}} + \frac{(n-1)(2n-1)b^3}{6n^3a^{3n-1}} - \frac{(n-1)(2n-1)(3n-1)b^4}{24n^4a^{4n-1}} + \text{etc.}, \end{aligned}$$

it will be having done the multiplication  $na^{n-1} + \alpha b + \beta b^2 + \gamma b^3 + \text{etc.}$

$$\begin{aligned} b = & b - \frac{(n-1)bb}{2na^n} + \frac{(n-1)(2n-1)b^3}{6n^2a^{2n}} - \frac{(n-1)(2n-1)(3n-1)b^4}{24n^3a^{3n}} + \text{etc.} \\ & + \frac{\alpha b^2}{na^{n-1}} - \frac{(n-1)\alpha b^3}{2n^2a^{2n-1}} + \frac{(n-1)(2n-1)\alpha b^4}{6n^3a^{3n-1}} \\ & + \frac{\beta b^3}{na^{n-1}} - \frac{(n-1)\beta b^4}{2n^2a^{2n-1}} \\ & + \frac{\gamma b^4}{na^{n-1}} \end{aligned}$$

Hence, the following determinations are deduced

$$\begin{aligned} \alpha &= \frac{n-1}{2a} \\ \beta &= \frac{(n-1)\alpha}{2na^n} - \frac{(n-1)(2n-1)}{6na^{n+1}} = -\frac{(n-1)(n+1)}{12na^{n+1}} \\ \gamma &= \frac{(n-1)\beta}{2na^n} - \frac{(n-1)(2n-1)\alpha}{6nna^{2n}} + \frac{(n-1)(2n-1)(3n-1)}{24n^2a^{2n+1}} = \frac{(n-1)(n+1)}{24na^{2n+1}}. \end{aligned}$$

Therefore, the fraction to be added to the already found root  $a$  will be

$$\frac{b}{na^{n-1} + \frac{(n-1)b}{2a} - \frac{(n-1)bb}{12na^{n+1}} + \frac{(n-1)b^3}{24na^{2n+1}} - \text{etc.}}$$

§238 Therefore, if the square root of the number  $N$  is to be extracted and already the approximate root was found  $= a$  together with the residue  $= b$ , to the found root one additionally has to add the quotient, which arises, if the residue  $b$  is divided by

$$2a + \frac{b}{2a} - \frac{bb}{8a^3} + \frac{b^3}{16a^5} - \text{etc.}$$

But if the cube root must be extracted, then the residue must be divided by

$$3a^2 + \frac{b}{a} - \frac{2bb}{9a^4} + \frac{b^3}{9a^7} - \text{etc.},$$

the use of which formulas we will show in these examples.

#### EXAMPLE 1

*Extract the square root of the number 200.*

Put  $N = 200$  and because the closest square is 196, it will be  $a = 14$  and the residue  $b = 4$ , which therefore must be divided by

$$28 + \frac{1}{7} - \frac{1}{7 \cdot 196} + \frac{1}{7 \cdot 196 \cdot 98},$$

and therefore the divisor will be  $= 28.142135$ ; if 4 is divided by it, one will obtain a decimal fraction to be added to 14, which will correct up to 10 figures and further.

#### EXAMPLE 2

*To extract the cube root of the number  $N = 10$ .*

The closest cube is 8 and the residue is  $= 2$ , whence  $a = 2$  and  $b = 2$  and the divisor  $= 12 + 1 - \frac{1}{18} = 12.9444$ . Hence, the cube root in question will approximately be  $= 2 \frac{2}{12.9444} = 2 \frac{10000}{64722}$ .

§239 The series found for the root can also be considered as a recurring series, arising from a certain fraction. For, this way many terms of the series will be reduced to a lot less, which constitute the numerator and the denominator of the fraction. So, having paid a little attention, one will see that it will be approximately

$$(a + b)^n = a^n \cdot \frac{a + \frac{n+1}{2}b}{a - \frac{n-1}{2}b}$$

and even closer

$$(a + b)^n = a^n \cdot \frac{aa + \frac{n+2}{2}ab + \frac{(n+1)(n+2)}{12}bb}{aa - \frac{n-2}{2}ab + \frac{(n-1)(n-2)}{12}bb}$$

In similar manner by introducing several terms even more accurate fractions can be obtained:

$$(a + b)^n = a^n \cdot \frac{a^3 + \frac{n+3}{2}a^2b + \frac{(n+3)(n+2)}{10}ab^2 + \frac{(n+3)(n+2)(n+1)}{120}b^3}{a^3 - \frac{n-3}{2}a^2b + \frac{(n-3)(n-2)}{10}ab^2 - \frac{(n-3)(n-2)(n-1)}{120}b^3}$$

Even a more general form of this kind be exhibited, to express which conveniently let be

$$\begin{array}{ll} A = \frac{m(n+m)}{1 \cdot 2m} & \mathfrak{A} = \frac{m(n-m)}{1 \cdot 2m} \\ B = \frac{(m-1)(n+m-1)}{2(2m-1)} A & \mathfrak{B} = \frac{(m-1)(n-m+1)}{2(2m-1)} \mathfrak{A} \\ C = \frac{(m-2)(n+m-2)}{3(2m-2)} B & \mathfrak{C} = \frac{(m-2)(n-m+2)}{3(2m-2)} \mathfrak{B} \\ D = \frac{(m-3)(n+m-3)}{4(2m-3)} C & \mathfrak{D} = \frac{(m-3)(n-m+3)}{4(2m-3)} \mathfrak{C} \\ \text{etc.} & \text{etc.} \end{array}$$

But having determined these values it will be

$$(a + b)^n = a^n \cdot \frac{a^m + Aa^{m-1}b + \mathfrak{B}a^{m-2}b^2 + Ca^{m-3}b^3 + \text{etc.}}{a^m - \mathfrak{A}a^{m-1}b + \mathfrak{B}a^{m-2}b^2 - \mathfrak{C}a^{m-3}b^3 + \text{etc.}}$$

§240 Therefore, if here for  $n$  a fractional number is substituted, these formulas will be very suitable for the extraction of roots. So if any root of power  $n$  must be extracted of the form  $a^n + b$ , the following formulas can be used

$$(a^n + b)^{\frac{1}{n}} = a \cdot \frac{2na^n + (n+1)b}{2na^n + (n-1)b}$$

$$(a^n + b)^{\frac{1}{n}} = a \cdot \frac{12n^2a^{2n} + 6n(2n+1)a^nb + (2n+1)(n+1)bb}{12n^2a^{2n} + 6n(2n-1)a^nb + (2n-1)(n-1)bb}$$

But if one puts  $a^n + b = N$  that it is  $a^n = N - b$ , it will be

$$(a^n + b)^{\frac{1}{n}} = a \cdot \frac{2nN - (n-1)b}{2nN - (n+1)b}$$

$$(a^n + b)^{\frac{1}{n}} a \cdot \frac{12n^2N^2 - 6n(2n-1)Nb + (2n-1)(n-1)bb}{12n^2N^2 - 6n(2n+1)Nb + (2n+1)(n+1)bb}$$

**§241** Therefore, the general formula for finding the root of any equation in equation, which consist of several terms, have the same use, which the usual rule of a binomial yields for the resolution of the pure equations  $x^n = c$ , and therefore in this case goes over into that rule itself. But if the equation was affected or even transcendental, our general expression is always used with the same success and yields an infinite series, which exhibits the value of the root. Therefore, since the greatest power of this formula consists in this task, let us show its use a little more diligently. Therefore, let this affected equations consisting of three terms be propounded

$$x^n + cx = N$$

while  $c$  and  $N$  denote any given quantities. Put  $x^n + cx - N = y$ ; it will be  $dy = (nx^{n-1} + c)dx$  and hence it will be  $p = \frac{1}{nx^{n-1} + c}$ ; then it is

$$dp = -\frac{n(n-1)x^{n-2}dx}{(nx^{n-1} + c)^2} \quad \text{and} \quad q = -\frac{n(n-1)x^{n-2}}{(nx^{n-1} + c)^3}$$

In similar manner because of  $r = \frac{dq}{dy}$ ,  $s = \frac{dr}{dy}$  etc. one will find

$$r = \frac{n^2(n-1)(2n-1)x^{2n-4} - n(n-1)(n-2)cx^{n-3}}{(nx^{n-1} + c)^5}$$

$$s = \frac{-n^3(n-1)(2n-1)(3n-1)x^{3n-6} + 4n^2(n-1)(n-2)(2n-1)cx^{2n-5} - n(n-1)(n-2)(n-3)c^2x^{n-4}}{(nx^{n-1} + c)^7}$$

$$t = \frac{\left\{ n^4(n-1)(2n-1)(3n-1)(4n-1)x^{4n-8} - n^3(n-1)(n-2)(2n-1)(29n-11)cx^{3n-7} \right\} + n^2(n-1)(n-2)(2n-1)(11n-29)c^2x^{2n-6} - n(n-1)(n-2)(n-3)(n-4)c^3x^{n-5}}{(nx^{n-1} + c)^9}$$

etc.

Having found these values the root of the propounded equation will be

$$f = x - py + \frac{1}{2}qyy - \frac{1}{6}ry^3 + \frac{1}{24}sy^4 - \frac{1}{120}ty^5 + \text{etc.};$$

for, whatever is substituted for  $x$ , whence at the same time the letters  $y, p, q, r$  etc, obtain determined values, the sum of the series will become equal to the value of one single root.

#### EXAMPLE 1

Let this equation be propounded  $x^3 + 2x = 2$ .

It will be  $c = 2, N = 2$  and  $n = 3$  and  $y = x^3 + 2x - 2$ . Put  $x = 1$ ; it will be  $y = 1$  and

$$p = \frac{1}{5}, \quad q = -\frac{6}{5^3}, \quad r = \frac{78}{5^5}, \quad s = -\frac{16 \cdot 90}{5^7} \quad \text{etc.}$$

and the root of the equation will be

$$f = -\frac{1}{5} - \frac{3}{5^3} - \frac{13}{5^5} - \frac{60}{5^7} - \text{etc.} = 0.771072.$$

Now put  $x = 0.77$ , and since it is  $y = x^3 + 2x - 2$ ,

$$p = \frac{1}{3xx + 2}, \quad q = -6p^3x, \quad r = 90xxp^5 - 12p^5$$

and

$$s = -2160p^7x^3 + 720p^7x,$$

by using logarithms one will have

$\log x^1$	$=$	9.8864907	$x^1$	$=$	0.77
$\log x^2$	$=$	9.7729814	$x^2$	$=$	0.5929
$\log x^3$	$=$	9.6594721	$x^3$	$=$	0.456533
			$2x$	$=$	1.54
			$x^3 + 2x$	$=$	1.996533
Therefore			$y$	$=$	-0.003467

And furthermore

$$\begin{array}{rcl}
 \log(-y) & = & 7.5399538 \\
 \log p & = & 9.4226575 \\
 \log(-py) & = & 6.9626113 \\
 \log p^3 & = & 8.2679725 \\
 \log x & = & 9.8864907 \\
 \log 3 & = & 0.4771213 \\
 \log y^2 & = & 5.0799076 \\
 \log(-\frac{1}{2}qyy) & = & 3.7114921 \\
 3xx + 2 & = & 3.7787 \\
 \log(3xx + 2) & = & 0.5773424 \\
 -py & = & 0.000917511 \\
 -\frac{1}{2}qyy & = & 0.000000514
 \end{array}$$

Therefore, the root is  $f = 0.770916997$ , which hardly deviates from the truth in the last figure.

#### EXAMPLE 2

Let the equation  $x^4 - 2xx + 4x = 8$  be propounded.

Put  $y = x^4 - 2xx + 4x - 8$ ; it will be  $dy = 4dx(x^3 - x + 1)$ ,

$$p = \frac{1}{4(x^3 - x + 1)}, \quad \frac{dp}{dx} = \frac{-3xx + 1}{4(x^3 - x + 1)^2}.$$

Therefore,

$$q = \frac{-3xx + 1}{16(x^3 - x + 1)^3}, \quad \frac{dq}{dx} = \frac{21x^4 - 12xx - 6x + 3}{16(x^3 - x + 1)^4} \quad \text{and} \quad r = \frac{21x^4 - 12xx - 6x + 3}{64(x^3 - x + 1)^5} \quad \text{etc.,}$$

from which the root of the propounded equation will be

$$f = x - \frac{y}{4(x^3 - x + 1)} - \frac{(3xx - 1)yy}{32(x^3 - x + 1)^3} - \frac{(7x^4 - 4xx - 2x + 1)y^3}{128(x^3 - x + 1)^5} - \text{etc.}$$

Therefore, it is necessary to attribute a suitable value to  $x$ , that series becomes convergent. But at first it is perspicuous, if such a value would be attributed to  $x$ , that it would be  $x^3 - x + 1 = 0$ , that then all terms of there series except the first would become infinite and nothing can be concluded from this. Therefore,

it is convenient to assign such a value of  $x$  that both  $y$  becomes small and  $x^3 - x + 1$  not very large. Let  $x = 1$ ; it will be  $y = -5$  and

$$f = 1 + \frac{5}{4} - \frac{25}{16} + \frac{125}{64} - \text{etc.};$$

because the three terms  $\frac{5}{4} - \frac{25}{16} + \frac{125}{64}$  agree with the terms of the geometric progression, whose sum is  $\frac{5}{9}$ , it will approximately be  $f = \frac{14}{9}$ . Therefore, let us put  $x = \frac{3}{2}$ ; it will be

$$y = -\frac{23}{16} \quad \text{and} \quad x^3 - x + 1 = \frac{23}{8},$$

whence it is

$$f = \frac{3}{2} + \frac{1}{8} - \frac{1}{64} + \frac{391}{256 \cdot 529} - \text{etc.} = 1.61.$$

Now put  $x = 1.61$ ; it will be

$\log x$	= 0.2068259	$x$	= 1.61	let $x^3 - x + 1 = z$
$\log x^2$	= 0.4136518	$x^2$	= 2.5921	
$\log x^3$	= 0.6204777	$x^3$	= 4.173281	
$\log x^4$	= 0.8273036	$x^4$	= 6.718983	
hence				
$\log(-y)$	= 8.4016934	$y$	= -0.025217	
$\log z$	= 0.5518502	$z$	= 3.563281	
$\log \frac{-y}{z}$	= 7.8498432			
$\log 4$	= 0.6020600			
$\log \frac{-y}{4z}$	= 7.2477832	$\frac{-y}{4z}$	= 0.0017692	
$\log(3xx - 1)$	= 0.8309926	$3xx - 1$	= 6.7763	
$\log y^2$	= 6.8033868			
	7.6343794			
$\log z^3$	= 1.6555506			
	5.9788288			
$\log 32$	= 1.5051500	$\frac{(3xx - 1)^2 y^2}{32z^3}$	= 0.00002976	
	= 4.4736788			



Therefore  $f = 1.6117662$ .

§242 This method to find the roots of equations approximately extends to transcendental quantities the same way. Let us find the number  $x$ , whose logarithm to the number itself has a given ratio as 1 to  $n$ , and one will have this equation  $x - n \log x$ ; but let  $k$  be the modulus of these logarithms, such that these logarithms are obtained, if the hyperbolic logarithms are multiplied by  $k$ ; it will be  $d. \ln x = \frac{kdx}{x}$ . Therefore, put  $x - n \log x = y$  and let  $f$  be the value of  $f$  in question which renders  $x = n \log x$ . Therefore, because it is  $y = x - n \log x$ , it will be

$$dy = dx - \frac{kndx}{x} = \frac{dx(x - kn)}{x}$$

and

$$\frac{dx}{dy} = p = \frac{x}{x - kn}, \quad \text{whence} \quad dp = -\frac{kndx}{(x - kn)^2},$$

therefore

$$\begin{aligned} \frac{dp}{dy} = q &= \frac{-knx}{(x - kn)^3}, & dq &= \frac{2knxdx + k^2n^2dx}{(x - kn)^4} \\ \frac{dq}{dy} = r &= \frac{knx(2x + kn)}{(x - kn)^5}. \end{aligned}$$

Hence, it will be

$$f = x - \frac{xy}{x - kn} - \frac{kxxy}{2(x - kn)^2} - \frac{kxxy^2(2x + kn)}{6(x - kn)^3} - \text{etc.}$$

Below [§ 272] we will show that this problem only admits a solution, if  $kn > e$  while  $e$  is the number, whose hyperbolic logarithm is = 1, or it must be  $kn > 2.7182818$ .

#### EXAMPLE

*A number except 10 shall be found, whose tabulated logarithm becomes equal to the tenth part of the number itself.*

Since the question is about tabulated logarithms, it will be  $k = 0.43429448190325$  and because of  $n = 10$  one will have  $kn = 4.3429448190325$ . Now, having put  $x = 1$  it will be  $y = 1$  and it will become

$$f = 1 + \frac{1}{3.3429} + \frac{2.1714724}{(3.3429)^3} - \text{etc.}$$

and so it will approximately be  $f = 1.37$ . Therefore, set  $x = 1.37$ ; it will be  $\log x = 0.136720567156406$  and because of  $y = x - 10 \log x$  it will be

$$y = 0.00279432843594 \quad \text{and} \quad -x + kn = 2.9729448190325.$$

Therefore, let

$$\begin{aligned} \log x &= 0.1367205 \\ \log y &= \frac{7.4462773}{7.5829978} \\ \log(kn - x) &= \frac{0.4731866}{7.1098112} \quad \frac{-xy}{x - kn} = 0.00128769 \end{aligned}$$

Further, because the third term is  $-\frac{knxyy}{2(x-kn)^2} = \frac{kny}{2(x-kn)^2} \cdot \frac{-xy}{x-kn}$ , it will be

$$\begin{aligned} \log \frac{-xy}{x - kn} &= 7.1098112 \\ \log y &= 7.4462773 \\ \log kn &= 0.6377842 \\ \hline &5.1938727 \\ \log(kn - x)^2 &= \frac{0.9463732}{4.2474995} \\ \log 2 &= 0.3010300 \\ \log \text{third term} &= 3.9464695 \\ \text{I. term } x &= 1.37 \\ \text{II. term} &= 0.00128769 \\ \text{III. term} &= 0.00000088 \\ \hline f &= 1.37128857 \\ \log f &= 0.137128857 \end{aligned}$$

§243 If the equation was an exponential equation, it can be reduced to any logarithmic one; so, if the value of  $x$  is in question, that it is  $x^x = a$ , it will be  $x \ln x = \ln a$ . Hence, having put  $y = x \ln x - \ln a$  it will be

$$dy = dx \ln x + dx \quad \text{and} \quad \frac{dx}{dy} = p = \frac{1}{1 + \ln x}$$

and then

$$dp = \frac{-dx}{x(1 + \ln x)^2} \quad \text{and} \quad \frac{dp}{dy} = q = \frac{-1}{x(1 + \ln x)^3},$$

$$dq = \frac{dx}{xx(1 + \ln x)^3} + \frac{3dx}{xx(1 + \ln x)^4} \quad \text{and hence} \quad \frac{dq}{dy} = r = \frac{1}{xx(1 + \ln x)^4} + \frac{3}{xx(1 + \ln x)^5};$$

further, it will be

$$dr = \frac{-2dx}{x^3(1 + \ln x)^4} - \frac{10dx}{x^3(1 + \ln x)^5} - \frac{15dx}{x^3(1 + \ln x)^6},$$

therefore

$$s = \frac{-2}{x^3(1 + \ln x)^5} - \frac{10}{x^3(1 + \ln x)^6} - \frac{15}{x^3(1 + \ln x)^7}$$

and

$$t = \frac{6}{x^4(1 + \ln x)^6} + \frac{40}{x^4(1 + \ln x)^7} + \frac{105}{x^4(1 + \ln x)^8} + \frac{105}{x^4(1 + \ln x)^9},$$

$$u = \frac{-24}{x^5(1 + \ln x)^7} - \frac{196}{x^5(1 + \ln x)^6} - \frac{700}{x^5(1 + \ln x)^9} - \frac{1260}{x^5(1 + \ln x)^{10}} - \frac{945}{x^5(1 + \ln x)^{11}}.$$

Therefore, if the true value of  $x$  is  $= f$ , such that  $f^f = a$ , it will be

$$f = x - \frac{y}{1 + \ln x} - \frac{yy}{2x(1 + \ln x)^3} - \frac{y^3}{2xx(1 + \ln x)^5} - \frac{5y^4}{8x^3(1 + \ln x)^7} - \frac{7y^5}{8x^4(1 + \ln x)^9}$$

$$- \frac{y^3}{6x^2(1 + \ln x)^4} - \frac{5y^4}{12x^3(1 + \ln x)^6} - \frac{7y^5}{8x^4(1 + \ln x)^8}$$

$$- \frac{y^4}{12x^3(1 + \ln x)^5} - \frac{y^5}{3x^4(1 + \ln x)^7}$$

$$- \frac{y^5}{20x^4(1 + \ln x)^6}$$

etc.

Therefore, this expression continued to infinity, whatever value is set for  $x$ , having taken  $y = x \ln x - \ln a$  will give the true value of  $f$ . So, if one puts  $x = 1$ , it will be  $y = -\ln a$  and

$$f = 1 + \ln a - \frac{(\ln a)^2}{2} + \frac{2(\ln a)^3}{3} - \frac{9(\ln a)^4}{8} + \frac{32(\ln a)^5}{15} - \frac{625(\ln a)^6}{144} - \text{etc.},$$

where it is to be noted that  $\ln a$  is the hyperbolic logarithm of  $a$ .

#### EXAMPLE

Find the number  $f$  that it is  $f^f = 100$ .

Because it is

$$a = 100 \quad \text{and} \quad y = x \ln x - \ln a = x \ln x - \ln 100,$$

since it is clear that it is  $f > 3$  and  $< 4$ , put  $x = \frac{7}{2}$  and it will be

$$\begin{aligned} \log x &= 1.25276296849 \\ x \log x &= 4.38467038972 \\ \log 100 &= 4.60517018599 \\ y &= \frac{-0.22049979627}{1 + \log x} \\ 1 + \log x &= 2.25276296849. \end{aligned}$$

Hence, by using ordinary logarithms it will be

$$\begin{aligned} \log(-y) &= 9.3434083 \\ \log(1 + \log x) &= 0.3527156 \\ \hline &8.9906927 \\ \log y^2 &= 8.6868166 \\ 3 \log(1 + \log x) &= 1.0581468 \\ \hline &7.6286698 \\ \log 2x = \log 7 &= 0.8450980 \\ \hline &6.7835718 \end{aligned} \quad \begin{aligned} \frac{-y}{1 + \log x} &= 0.0978797 \\ \frac{y^2}{2x(1 + \log x)^3} &= 0.0006075. \end{aligned}$$

Therefore, it will approximately be  $f = 3.5972722$ ;

but having additionally taken the following terms, it will be  $f = 3.5972852$ .

§244 But moreover differential calculus has an extraordinary use in the resolution of equations, if a certain relation, which intercedes between the roots, was known. Let the equation  $y = 0$  be propounded, in which  $y$  shall be any function of  $x$ . If now, for the sake of an example, it is known that the roots of this equation differ by the given quantity  $a$ , these two roots will easily be found the following way. Let  $x$  denote the smaller of these two roots; the larger will be  $= x + a$ ; hence, because the function  $y$  vanishes, if  $x$  denotes any of the roots of the equation  $y = 0$ , it will also vanish, if instead of  $x$  one puts  $x + a$ . Therefore, it will be

$$0 = y + \frac{ady}{dx} + \frac{a^2ddy}{2dx^2} + \frac{a^3d^3y}{6dx^3} + \text{etc.}$$

Hence, because it is  $y = 0$ , it will also be

$$0 = \frac{ady}{dx} + \frac{a^2ddy}{2dx^2} + \frac{a^3d^3y}{6dx^3} + \text{etc.}$$

which two equations taken at the same time by the method of elimination will give the value of the root  $x$ , which is smaller than the other root by the quantity  $a$ .

#### EXAMPLE

Let this equation  $x^5 - 24x^4 + 49xx - 36 = 0$  be propounded, which is known from anywhere to have two roots differing by the unity.

Having put  $y = x^5 - 24x^3 + 49xx - 36$  it will be

$$\begin{aligned} \frac{dy}{dx} &= 5x^4 - 72x^2 + 98x \\ \frac{ddy}{2dx^2} &= 10x^3 - 72x + 49 \\ \frac{d^3y}{6dx^3} &= 10x^2 - 24 \\ \frac{d^4y}{24dx^4} &= 5x \\ \frac{d^5y}{120dx^5} &= 1. \end{aligned}$$

But because of  $a = 1$  it will be

$$A \dots 5x^4 + 10x^3 - 62x^2 + 31x + 26 = 0.$$

But it is

$$B \dots x^5 - 24x^3 + 49xx - 36 = 0.$$

Multiply the superior by  $x$  and the inferior by 5 and subtract the one from the other and it will remain

$$10x^4 + 58x^3 - 214x^2 + 26x + 180 = 0$$

or

$$C \dots 5x^4 + 229x^3 - 107x^2 + 13x + 90 = 0,$$

from which the first  $A$  subtracted will give

$D \dots$	$19x^3$	$-$	$45x^2$	$-$	$18x$	$+$	$64$	$=$	$0.$
$D \cdot 5x \dots$	$95x^4$	$-$	$225x^3$	$-$	$90x^2$	$+$	$320x$	$=$	$0.$
$A \cdot 19 \dots$	$95x^4$	$+$	$190x^3$	$-$	$1178x^2$	$+$	$589x$	$+$	$494 = 0.$
$E \dots$			$415x^3$	$-$	$1088x^2$	$+$	$269x$	$+$	$494 = 0.$
$D \cdot 415 \dots$			$7885x^3$	$-$	$18675x^2$	$-$	$7470x$	$+$	$26560 = 0.$
$E \cdot 19 \dots$			$7885x^3$	$-$	$20672x^2$	$+$	$5111x$	$+$	$8386 = 0.$
$F \dots$					$1997x^2$	$-$	$12581x$	$+$	$17174 = 0.$

And

$$\begin{array}{rcccccccc}
 D \cdot 247 \dots & 4693x^3 & - & 11115x^2 & - & 4446x & + & 15808 & = & 0 \\
 D \cdot 32 \dots & 13280x^3 & - & 34816x^2 & + & 8608x & + & 15808 & = & 0 \\
 \hline
 & 8587x^3 & - & 23701x^2 & + & 13054x & = & 0 & & \\
 \hline
 G \dots & & & 8587x^2 & - & 23701x & + & 13054 & = & 0 \\
 F \cdot 8587 \dots & & & 17148239x^2 & - & 108033047x & + & 147473138 & = & 0 \\
 G \cdot 1997 \dots & & & 17148239x^2 & - & 47330897x & + & 26068838 & = & 0 \\
 \hline
 & & & & & 60702150x & - & 121404300 & = & 0.
 \end{array}$$

From this equation it follows  $x = 2$  and therefore also  $x = 3$  will be a root of the equation, both of which values indeed satisfy the equation.

§245 But this operation can be done without use of differential calculus, because the same equation, which the differential calculus yielded, arises, if in the propounded equation one puts  $x + 1$  instead of  $x$ . Furthermore, this method of elimination is too laborious, and if the equations would be of higher degree, the labor would be simply too much to handle; and this holds even more for transcendental equations. But if we put that two roots of the propounded equation  $y = 0$  are equal to each other, then because of  $x = a$  the differential equations goes over into this one  $\frac{dy}{dx} = 0$ . Therefore, if any equation  $y = 0$  had two equal roots, it will be  $\frac{dy}{dx} = 0$  and these two roots taken together will yield the value of  $x$ , to which these two roots are equal. Hence, vice versa, if the two equations  $y = 0$  and  $\frac{dy}{dx} = 0$  have a common root, it will be a double root of the equation  $y = 0$ . But this happens, if, after the quantity  $x$  by means of these two equations  $y = 0$  and  $\frac{dy}{dx} = 0$  was completely eliminated, one gets to an identical equation. So if the equation

$$x^3 - 2xx - 4x + 8 \tag{1}$$

was propounded, it will also be  $3xx - 4x - 4 = 0$ , whose double added to the first gives

$$x^3 + 4xx - 12x = 0 \quad \text{or} \quad xx + 4x - 12 = 0,$$

whose triple is

$$\begin{array}{r}
3xx + 12x - 36 = 0 \\
\text{subtract } 3xx - 4x - 4 = 0 \\
\hline
16x - 32 = 0 \\
x - 2 = 0.
\end{array}$$

Therefore, because  $x = 2$  arose, substitute this value in one of the preceding  $3xx - 4 - 4 = 0$  and the identical equation  $12 - 8 - 4 = 0$  will arise, whence one concludes that the propounded equation  $x^3 - 2xx - 4x + 8 = 0$  has to equal roots, namely  $x = 2$ .

§246 If one therefore has an algebraic equation of no matter how many dimensions

$$x^n + Ax^{n-1} + Bx^{n-2} + Cx^{n-3} + Dx^{n-4} + \text{etc.} = 0,$$

which has two equal roots, it will also be

$$nx^{n-1} + (n-1)Ax^{n-2} + (n-2)Bx^{n-2} + (n-3)Cx^{n-4} + (n-4)Dx^{n-5} + \text{etc.} = 0.$$

This double root of that equation will at the same time be a root of the first equation, of course. Multiply the first equation by  $n$  and from it subtract the second multiplied by  $x$  and this new equation will arise

$$Ax^{n-1} + 2Bx^{n-2} + 3Cx^{n-3} + 4Dx^{n-4} + \text{etc.} = 0.$$

Now add the first multiplied by  $a$  and the latter multiplied by  $b$ ; it will be

$$ax^n + (a+b)Ax^{n-1} + (a+2b)Bx^{n-2} + (a+3b)Cx^{n-3} + \text{etc.} = 0,$$

which equation combined with the propounded itself will show equal roots, if the propounded one has some. Therefore, because the quantities  $a$  and  $b$  can be taken ad libitum, the coefficients  $a$ ,  $a+b$ ,  $a+2b$  etc. represent any arithmetic progression. Therefore, if any equation has to equal roots, they will be found, if the single terms of the propounded equation are multiplied by terms of a certain arithmetic progression, respectively; for, the new equation resulting this way will also contain the root, which is contained twice in the propounded one. So, if the the terms of the equation



$$x^n + Ax^{n-1} + Bx^{n-2} + Cx^{n-3} + Dx^{n-4} + \text{etc.}$$

are multiplied by this arithmetic progression

$$a, \quad a + b, \quad a + 2b, \quad a + 3b, \quad a + 4b \quad \text{etc.};$$

this new equation will arise

$$ax^n + (a + b)Ax^{n-1} + (a + 2b)Bx^{n-2} + (a + 3b)Cx^{n-3} + (a + 4b)Dx^{n-4} + \text{etc.} = 0,$$

which combined with the latter will show the equal roots. And this is the sufficiently known rule to find equal roots of any equation.

§247 If the equation  $y = 0$  has three equal roots, it will not only be  $\frac{dy}{dx} = 0$ , but it will also be  $\frac{ddy}{dx^2} = 0$ , if for  $x$  one sets the value of the root, which is in the equation  $y = 0$  trice. To show this let us put that the equation  $y = 0$  has three roots  $x, x + a, x + b$  etc., of which the first differs to the other ones by  $a$  and  $b$ , respectively; and since  $y$  vanishes, if instead of  $x$  one writes  $x + a$  or  $x + b$ , it will be

$$\begin{aligned} y &= 0 \\ y + \frac{ady}{dx} + \frac{a^2ddy}{2dx^2} + \frac{a^3d^3y}{6dx^3} + \frac{a^4d^4y}{24dx^4} + \text{etc.} &= 0 \\ y + \frac{bdy}{dx} + \frac{b^2ddy}{2dx^2} + \frac{b^3d^3y}{6dx^3} + \frac{b^4d^4y}{24dx^4} + \text{etc.} &= 0; \end{aligned}$$

if from the two latter ones the first is subtracted, it will be

$$\begin{aligned} \frac{dy}{dx} + \frac{addy}{2dx^2} + \frac{a^2d^3y}{6dx^3} + \frac{a^3d^4y}{24dx^4} + \text{etc.} &= 0 \\ \frac{dy}{dx} + \frac{bdy}{2dx^2} + \frac{b^2d^3y}{6dx^3} + \frac{b^3d^4y}{24dx^4} + \text{etc.} &= 0 \end{aligned}$$

Also subtract these from each other and having divided by  $a - b$  it will be

$$\frac{ddy}{2dx^2} + \frac{(a + b)d^3y}{6dx^3} + \frac{(aa + ab + bb)d^4y}{24dx^4} + \text{etc.} = 0.$$

Now put  $a = 0$  and  $b = 0$  such that these three roots are equal to each other, and because of the vanishing terms it will be

$$y = 0, \quad \frac{dy}{dx} = 0 \quad \text{and} \quad \frac{ddy}{dx^2} = 0.$$

**§248** Therefore, if the equation  $y = 0$  has three equal roots, say  $f, f, f$ , then this quantity  $f$  will also be a root not only of this equation  $\frac{dy}{dx} = 0$ , but also of this one  $\frac{ddy}{dx^2} = 0$ . Hence, it is manifest, because  $f$  is the common root of the equation  $\frac{dy}{dx} = 0$  and its differential  $\frac{ddy}{dx^2} = 0$ , that it has to be contained twice in the equation  $\frac{dy}{dx} = 0$  by means of that what we showed before about two equal roots of equations. Hence, if the equation

$$x^n + Ax^{n-1} + Bx^{n-2} + Cx^{n-3} + Dx^{n-4} + \text{etc.} = 0$$

contains three equal roots  $f, f, f$ , if its terms are multiplied by the terms of a certain arithmetic progression, then the resulting equation will have two equal roots  $f$  and  $f$ ; therefore, it can be multiplied by an arithmetic progression again, that an equation arises containing the root  $f$  once. Therefore, one will obtain three equations having the common roots  $f$ , from whose combination this root will easily be found. For, if arithmetic progressions of such a kind are chosen, whose either first or last terms shall be  $= 0$ , then an equation of one degree lower will arise and so the elimination will become even easier.

**§249** In similar manner it will be shown, if the equation  $y = 0$  has four equal roots  $f, f, f, f$ , that than for  $x = f$  will not only become  $y = 0$ ,  $\frac{dy}{dx} = 0$  and  $\frac{ddy}{dx^2} = 0$ , but it will also be  $\frac{d^3y}{dx^3} = 0$ . As the equation  $y = 0$  contains the root  $x = f$  four times, so the equation  $\frac{dy}{dx}$  will contain the same thrice, the equation  $\frac{ddy}{dx^2} = 0$  twice and the equation  $\frac{d^3y}{dx^3} = 0$  once. This will also be seen easier, if we consider that the function  $y$  in this case has to have a form of this kind  $(x - f)^4 X$ , where  $X$  denotes any function of  $x$ . Having assumed this form it will be

$$\frac{dy}{dx} = (x - f)^3 \left( 4X + \frac{(x - f)dX}{dx} \right)$$

and hence be divisible by  $(x - f)^3$ . Similarly further  $\frac{ddy}{dx^2}$  will have the factor  $(x - f)^2$  and  $\frac{d^3y}{dx^3}$  the factor  $x - f$ ; from this it is perspicuous, if the root  $f$  is

contained in the equation  $y = 0$ , it has to be contained in the equation  $\frac{dy}{dx} = 0$  trice, in the equation  $\frac{d^2y}{dx^2} = 0$  twice and in  $\frac{d^3y}{dx^3} = 0$  still once.